

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2007 (09.08.2007)

PCT

(10) International Publication Number
WO 2007/089470 A2

(51) International Patent Classification:

A61K 39/40 (2006.01)

(21) International Application Number:

PCT/US2007/001687

(22) International Filing Date: 23 January 2007 (23.01.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/763,023 27 January 2006 (27.01.2006) US

(71) Applicant (for all designated States except US): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BROWN, Martha, J.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **ANDERSON, Annaliesa, S.** [GB/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **COPE, Leslie, D.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **JANSEN, Kathrin, Ute** [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **MCNEELY, Tessie**

[US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **HARVEY, Barrett** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **DURR, Erberhard** [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **ERNST, Robin** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).


(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: ANTIGEN-BINDING PROTEINS TARGETING S. AUREUS ORF0657N



(57) Abstract: The present invention features antigen binding protein that bind an ORF0657n target region (SEQ ID NO: 1). ORF0657n is an *S. aureus* protein. ORF0657n target regions are provided by the mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 binding sites. In a lethal model challenge, mAb 2H2.BE11 and mAb 13C7.BC1 provided for increased survival against *S. aureus* infection. There was also protection demonstrated in an *ex vivo* model with either the IgG1 or the IgG2b form of mAb 2H2; and in a passive immunization murine indwelling catheter model using mAb 2H2.BE11.



WO 2007/089470 A2



GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE OF THE INVENTION**ANTIGEN-BINDING PROTEINS TARGETING *S. AUREUS* ORF0657n****RELATED APPLICATIONS**

The present application claims priority to U.S. Provisional Application No. 60/763,023, filed January 27, 2006, which is hereby incorporated by reference herein.

BACKGROUND OF THE INVENTION

The references cited throughout the present application are not admitted to be prior art to the claimed invention.

Staphylococcus aureus is a pathogen responsible for a wide range of diseases and conditions. Examples of diseases and conditions caused by *S. aureus* include bacteremia, infective endocarditis, folliculitis, furuncle, carbuncle, impetigo, bullous impetigo, cellulitis, botryomycosis, toxic shock syndrome, scalded skin syndrome, central nervous system infections, infective and inflammatory eye disease, osteomyelitis and other infections of joints and bones, and respiratory tract infections. (*The Staphylococci in Human Disease*, Crossley and Archer (eds.), Churchill Livingstone Inc. 1997.)

Immunological based strategies can be employed to control *S. aureus* infections and the spread of *S. aureus*. Immunological based strategies include passive and active immunization. Passive immunization employs immunoglobulins targeting *S. aureus*. Active immunization induces immune responses against *S. aureus*.

SUMMARY OF THE INVENTION

The present invention features antigen binding protein that bind an ORF0657n target region (SEQ ID NO: 1). ORF0657n is an *S. aureus* protein. ORF0657n target regions are provided by the mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 binding sites. In a lethal model challenge, mAb 2H2.BE11 and mAb 13C7.BC1 provided for increased survival against *S. aureus* infection. There was also protection demonstrated in an *ex vivo* model with either the IgG1 or the IgG2b form of mAb 2H2; and in a passive immunization murine indwelling catheter model using mAb 2H2.BE11.

Mouse hybridoma cell lines producing mAb 1G3.BD4, mAb 2H2.BE11; mAb 13C7.BC1, and mAb 13G11.BF3 were deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, in accordance with Budapest Treaty on September 30, 2005. The cells lines were designated: ATCC No. PTA-7124 (producing mAb 2H2.BE11), ATCC No. PTA-7125 (producing mAb 13C7.BC1), ATCC No. PTA-7126 (producing mAb 1G3.BD4), and ATCC No. PTA-7127 (producing mAb 13G11.BF3).

Thus, a first aspect of the present invention features an isolated antigen binding protein comprising a first variable region and a second variable region. The first and second variable regions

bind one or more target regions selected from the group consisting of: mAb 1G3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1 target region, and mAb 13G11.BF3 target region.

Reference to "isolated" indicates a different form than found in nature. The different form can be, for example, a different purity than found in nature and/or a structure that is not found in nature. A structure not found in nature includes recombinant structures where different regions are combined together, for example, humanized antibodies where one or more murine complementary determining regions is inserted onto a human framework scaffold or a murine antibody is resurfaced to resemble the surface residues of a human antibody, hybrid antibodies where one or more complementary determining regions from an antigen binding protein is inserted into a different framework scaffold, and antibodies derived from natural human sequences where genes coding for light and heavy variable domains were randomly combined together.

The isolated protein is preferably substantially free of serum proteins. A protein substantially free of serum proteins is present in an environment lacking most or all serum proteins.

A "variable region" has the structure of an antibody variable region from a heavy or light chain. Antibody heavy and light chain variable regions contain three complementary determining regions interspaced onto a framework. The complementary determining regions are primarily responsible for recognizing a particular epitope.

A target region is defined with respect to the ORF0657n region (SEQ ID NO: 1) bound by mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, or mAb 13G11.BF3. For example, the mAb 1G3.BD4 target region is the ORF0657n region to which mAb 1G3.BD4 binds.

A protein binding an identified target region competes with either mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, or mAb 13G11.BF3 for binding to the target region. For example, a protein competing with mAb 1G3.BD4 binding to ORF0657n binds to the mAb 1G3.BD4 target region.

A protein that competes with either the monoclonal antibody mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11 reduces binding of the monoclonal antibody to ORF0657n by at least about 20%, preferably at least about 50%, when excess and equal amounts of the competing protein and monoclonal antibody are employed.

Reference to "protein" indicates a contiguous amino acid sequence and does not provide a minimum or maximum size limitation. One or more amino acids present in the protein may contain a post-translational modification, such as glycosylation or disulfide bond formation.

A preferred antigen binding protein is a monoclonal antibody. Reference to a "monoclonal antibody" indicates a collection of antibodies having the same, or substantially the same, complementary determining region, and binding specificity. The variation in the antibodies is that which would occur if the antibodies were produced from the same construct(s).

Monoclonal antibodies can be produced, for example, from a particular hybridoma and from a recombinant cell containing one or more recombinant genes encoding the antibody. The antibody

may be encoded by more than one recombinant gene where, for example, one gene encodes the heavy chain and one gene encodes the light chain.

Another aspect of the present invention describes a nucleic acid containing a recombinant gene comprising a nucleotide sequence encoding an antibody variable region. The antibody variable region can bind a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region.

A recombinant gene contains recombinant nucleic acid encoding a protein along with regulatory elements for proper transcription and processing (which may include translational and post translational elements). The recombinant nucleic acid by virtue of its sequence and/or form does not occur in nature. Examples of recombinant nucleic acid include purified nucleic acid, two or more nucleic acid regions combined together providing a different nucleic acid than found in nature, and the absence of one or more nucleic acid regions (*e.g.*, upstream or downstream regions) that are naturally associated with each other.

Another aspect of the present invention describes a recombinant cell comprising one or more recombinant genes encoding an antibody variable region that binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region. Multiple recombinant genes are useful, for example, where one gene encodes an antibody heavy chain or fragment thereof containing the V_H region and another nucleic acid encodes an antibody light chain or fragment thereof containing the V_L region.

Another aspect of the present invention comprises a method of producing a protein comprising an antibody variable region. The method comprising the steps of: (a) growing a recombinant cell comprising recombinant nucleotide acid encoding for a protein under conditions wherein the protein is expressed; and (b) purifying the protein.

Another aspect of the present invention describes a pharmaceutical composition. The composition contains a therapeutically effective amount of an antigen binding protein and a pharmaceutically acceptable carrier.

A therapeutically effective amount is an amount sufficient to provide a useful therapeutic or prophylactic effect. For a patient infected with *S. aureus*, an effective amount is sufficient to achieve one or more of the following effects: reduce the ability of *S. aureus* to propagate in the patient or reduce the amount of *S. aureus* in the patient. For a patient not infected with *S. aureus*, an effective amount is sufficient to achieve one or more of the following: a reduced susceptibility to *S. aureus* infection or a reduced ability of the infecting bacterium to establish persistent infection for chronic disease.

Another aspect of the present invention describes a method of detecting the presence of an OFR0657n antigen in a solution or on a cell. The method involves providing a binding protein described herein to the solution or cell and measuring the ability of the binding protein to bind to the antigen in the solution or cell. Measurements can be quantitative or qualitative.

Reference to ORF0657n antigen includes full-length ORF0657n or a derivative thereof having an epitope that is recognized by mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11. Examples of derivatives include truncated versions; and full-length or truncated versions of ORF0657n containing one or more of the following amino acid alterations: one or more additions, one or more substitutions, and one or more deletions.

Another aspect of the present invention features a method of treating a patient against a *S. aureus* infection. The method comprises the step of administering to the patient an effective amount of an antigen binding protein described herein. The patient being treated may, or may not, be infected with *S. aureus*. Preferably, the patient is a human.

Another aspect of the present invention describes a cell line producing a protein that is either mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11, or that competes with either mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11 for binding to ORF0657n. Preferred cells lines are hybridomas, and recombinant cell lines containing recombinant nucleic acid encoding the protein.

Reference to open-ended terms such as "comprises" allows for additional elements or steps. Occasionally phrases such as "one or more" are used with or without open-ended terms to highlight the possibility of additional elements or steps.

Unless explicitly stated reference to terms such as "a" or "an" is not limited to one. For example, "a cell" does not exclude "cells". Occasionally phrases such as one or more are used to highlight the possible presence of a plurality.

Other features and advantages of the present invention are apparent from the additional descriptions provided herein including the different examples. The provided examples illustrate different components and methodology useful in practicing the present invention. The examples do not limit the claimed invention. Based on the present disclosure the skilled artisan can identify and employ other components and methodology useful for practicing the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the structure of an IgG molecule. "V_L" refers to a light chain variable region. "V_H" refers to a heavy chain variable region. "C_L" refers to a light chain constant region. "CH₁", "CH₂" and "CH₃" are heavy chain constant regions. Dashed lines indicate disulfide bonds.

Figure 2 illustrates a matrix outlining the reactivities of different monoclonal antibodies in a pair-wise binding study. The panel of monoclonal antibodies fell into three reactive areas by the BIAcore® method.

Figures 3A-3C: Groups of BALB/c mice (n = 20) were treated 20 hours prior to bacterial challenge with an i.p. injection of: ■, mAb 13C7.BC1; □, mAb 6G6.A8 (isotype control); or ○, PBS. Mice were challenged with *S. aureus* by i.v. injection and survival was monitored. Fig. 3A-

0.49 mg mAb 13C7.BC1; 0.45 mg mAb 6G6.A8; and 9.8×10^8 CFU *S. aureus* Becker. Fig. 3B- 0.49 mg mAb 13C7.BC1; 0.45 mg mAb 6G6.A8; and 9.6×10^8 CFU *S. aureus* Becker. Fig. 3C- 0.50 mg mAb 13C7.BC1; 0.45 mg mAb 6G6; and 9.9×10^8 CFU *S. aureus* Becker.

Figures 4A and 4B: Groups of BALB/c mice ($n = 20$) were treated 20 hours prior to bacterial challenge with an i.p. injection of: ■, mAb 13C7.BC1 (0.5 mg); □, mAb 6G6.A8 (isotype control) (0.5 mg); or ○, PBS (0.5 ml). Mice were challenged with *S. aureus* by i.v. injection and survival was monitored. Fig. 4A illustrates results with 2.09×10^8 CFU *S. aureus* UK58. Fig. 4B illustrates results with 2.15×10^8 *S. aureus* UK 58.

Figures 5A-5C: Groups of BALB/c mice ($n = 20$) were treated 20 hours prior to bacterial challenge with an i.p. injection of: ■, mAb 2H2.BE11, □, mAb 6G6.A8 (isotype control); ○, PBS. Mice were challenged with *S. aureus* by i.v. injection and survival was monitored. Fig. 5A- 0.43 mg mAb 2H2.BE11; 0.5 mg mAb 6G6.A8; and 9.8×10^8 CFU *S. aureus* Becker. Fig. 5B- 0.43 mg mAb 2H2.BE11; 0.5 mg mAb 6G6.A8; and 8.3×10^8 CFU *S. aureus* Becker. Fig. 5C- 0.43 mg mAb 2H2.BE11; 0.5 mg mAb 6G6.A8; and 9.3×10^8 CFU *S. aureus* Becker.

DETAILED DESCRIPTION OF THE INVENTION

ORF0657n is an *S. aureus* protein located at the *S. aureus* outer membrane. ORF0657n has been found to be well conserved in different strains of *S. aureus*. (Anderson *et al.*, International Publication No. WO 2005/009379, International Publication Date February 3, 2005.) Different ORF0657n derivatives can be used to produce a protective immune response against *S. aureus* infection. (Anderson *et al.*, International Publication No. WO 2005/009379, International Publication Date February 3, 2005.)

Due to their ability to recognize ORF0657n, the antigen binding proteins described herein can be used, for example, as a tool in the production, characterization, or study of ORF0657n based antigens. Antigen binding protein recognizing appropriate ORF0657n epitopes can also be used agent to treat *S. aureus* infection.

I. Antigen Binding Protein

Antigen binding proteins contain an antibody variable region providing for specific binding to an epitope. The antibody variable region can be present in, for example, a complete antibody, an antibody fragment, and a recombinant derivative of an antibody or antibody fragment.

Different classes of antibodies have different structures. Different antibody regions can be illustrated by reference to IgG (Figure 1). An IgG molecule contains four amino acid chains: two longer length heavy chains and two shorter light chains. The heavy and light chains each contain a constant region and a variable region. Within the variable regions are three hypervariable regions responsible for antigen specificity. (See, for example, Breitling *et al.*, Recombinant Antibodies, John Wiley & Sons, Inc. and Spektrum Akademischer Verlag, 1999; and Lewin, Genes IV, Oxford University Press and Cell Press, 1990.)

The hypervariable regions (also referred to as complementarity determining regions), are interposed between more conserved flanking regions (also referred to as framework regions). Amino acids associated with framework regions and complementarity determining regions can be numbered and aligned as described by Kabat *et al.*, Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services, 1991.

The two heavy chain carboxyl regions are constant regions joined by disulfide binding to produce an Fc region. The Fc region is important for providing antibody biological activity such as complement and macrophage activation. Each of the two heavy chains making up the Fc region extend into different Fab regions through a hinge region.

In higher vertebrates there are two classes of light chains and five classes of heavy chains. The light chains are either κ or λ . The heavy chains define the antibody class and are either α , δ , ϵ , γ , or μ . For example, IgG has a γ heavy chain. Subclasses also exist for different types of heavy chains such as human γ_1 , γ_2 , γ_3 , and γ_4 . Heavy chains impart a distinctive conformation to hinge and tail regions. (Lewin, Genes IV, Oxford University Press and Cell Press, 1990.)

Antibody fragments containing an antibody variable region include Fv, Fab, and Fab₂ regions. Each Fab region contains a light chain made up of a variable region and a constant region, and a heavy chain region containing a variable region and a constant region. A light chain is joined to a heavy chain by disulfide bonding through constant regions. The light and heavy chain variable regions of a Fab region provide for an Fv region that participates in antigen binding.

The antibody variable region can be present in a recombinant derivative. Examples of recombinant derivatives include single-chain antibodies, diabody, triabody, tetrabody, and miniantibody. (Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004.)

The antigen binding protein can contain one or more variable regions recognizing the same or different epitopes. (Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004.)

II. Generation of Antigen Binding Protein Directed to an Identified Target Region

Different antigen binding proteins directed to the mAb 1G3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1 target region, or mAb 13G11.BF3 target region can be generated starting with the respective monoclonal antibody. Alternatively, the epitope recognized by a binding protein can be used to select additional binding proteins.

The mAb 2H2.BE11 target region appears to be located at approximately amino acids 76-357 of ORF0657n. A polypeptide containing amino acids 76-357 of ORF0657n, or a full-length ORF0657n, can be used as a target antigen to select for antibodies. The target region of the generated antibodies can be determined.

A variety of techniques are available to select for a protein recognizing an antigen. Examples of such techniques include use of phage display technology and hybridoma production. Human antibodies can be produced using chimeric mice such as a XenoMouse or Trans-Chromo mouse.

(E.g., Azzazy *et al.*, *Clinical Biochemistry* 35:425-445, 2002, Berger *et al.*, *Am. J. Med. Sci.* 324(1):14-40, 2002.)

The monoclonal antibodies mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 contain variable regions recognizing ORF0675n. Additional binding proteins recognizing ORF0657n can be produced based on antibody variable regions. Additional binding proteins can, for example, be produced by modifying an existing monoclonal antibody and by using variable region sequence information. Protein construction and sequence manipulation can be performed using recombinant nucleic acid techniques.

The monoclonal antibodies mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 are murine antibodies. For human therapeutic applications, preferred binding proteins based on such mAb's are designed to reduce the potential generation of human anti-mouse antibodies recognizing the murine regions.

The potential generation of human anti-mouse antibodies can be reduced using techniques such as murine antibody humanization, de-immunization, and chimeric antibody production. (See, for example, O'Brien *et al.*, Humanization of Monoclonal Antibodies by CDR Grafting, p 81-100, From *Methods in Molecular Biology* Vol. 207: Recombinant antibodies for Cancer Therapy: Methods and Protocols (Eds. Welschhof and Krauss) Humana Press, Totowa, New Jersey, 2003; Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004; Gonzales *et al.*, *Tumor Biol.* 26:31-43, 2005, Presta, *Advanced Drug Delivery Reviews* 58:640-656, 2006, Tsurushita *et al.*, *Methods* 36:69-83, 2005, Roque *et al.*, *Biotechnol. Prog.* 20:639-654, 2004.)

Murine antibodies can be humanized using techniques such as grafting complementary determining regions into a framework region or resurfacing. Resurfacing (also known as veneering) involves modifying a variable region so the surface exposed regions are humanized.

Grafting complementary determining regions involves taking such regions or a portion of such regions from, for example, a murine source and inserting the regions into a human variable region framework. The human framework used for grafting can be selected based on sequence homology to the variable region (e.g., murine) from which the region was obtained. Essential framework residues associated with grafted complementary determining regions should also be provided in the new framework.

De-immunization involves altering potential linear T-cell epitopes present in the antibody. The epitopes can be identified based on a bioinformatics scan of known human HLA class I and/or class II epitopes. (Presta, *Advanced Drug Delivery Reviews* 58:640-656, 2006.)

A chimeric antibody contains a human constant region along with a variable region from a different organism, such as a mouse. The human constant region provides an Fc region.

Additional examples of alterations include providing a variable region in, for example, a single chain antibody, a diabody, a triabody, a tetrabody, and a miniantibody. (Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004.) The antigen binding protein can contain one or more variable

regions recognizing the same or different epitopes. (*Id.*) Additional embodiments of the present invention are directed to a single chain antibody, a diabody, a triabody, a tetrabody, or a miniantibody directed to the mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, or mAb 13G11.BF3 binding site.

III. Binding Protein Directed to the mAb 2H2.BE11 Target Region

As described in the Examples provided below, the mAb 2H2.BE11 target region was further characterized and the amino acids sequence of the variable regions was determined. The identified target region and the sequence information facilitate obtaining different binding proteins directed to the mAb 2H2.BE11 target region.

In an embodiment of the present invention, the binding protein binds to a polypeptide consisting of amino acids 76-357 of SEQ ID NO: 1. Preferably, the binding protein is either a human antibody, a humanized antibody, a de-immunized antibody, or chimeric antibody. Preferred antibodies are isolated antibodies and monoclonal antibodies.

The amino acids sequences of the mAb 2H2.BE11 variable regions are provided by SEQ ID NO: 20 (V_h) and SEQ ID NO: 21 (V_l). The complementary determining regions (CDR's) within V_h were identified at amino acids 36-45, 50-65, and 98-107. The CDR's within V_l were identified at amino acids 24-33, 49-55, and 88-96 of SEQ ID NO: 21.

In different embodiments directed to a V_h region, the binding protein binds the mAb 2H2.BE11 target region and comprises, consists, or consists essentially of: a first V_h CDR comprising, consisting, or consisting essentially of amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid; a second V_h CDR comprising, consisting, or consisting essentially of amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acid; and a third V_h CDR comprising, consisting, or consisting essentially of amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.

In different embodiments directed to a V_l region, the binding protein binds the mAb 2H2.BE11 target region and comprises, consists, or consists essentially of a first V_l CDR comprising, consisting, or consisting essentially of amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid; a second V_l CDR comprising, consisting, or consisting essentially of amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and a third V_l CDR comprising, consisting, or consisting essentially of amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.

Reference to "consisting essentially of" with respect to a variable region, CDR region, or antibody sequence, indicates the possible presence of one or more additional amino acids, where such amino acids do not significantly decrease binding to the target.

An amino acid difference can be an amino acid deletion, insertion, or substitution. In substituting amino acids to maintain activity, the substituted amino acids should have one or more similar properties such as approximately the same charge, size, polarity and/or hydrophobicity.

Preferably, an amino acid substitution is a conservative substitution. A conservative substitution replaces an amino acid with another amino acid having similar properties. Table 1 provides a list of groups of amino acids, where one member of the group is a conservative substitution for another member.

Table 1 : Conservative Substitutions

Ala, Val, Ile, Leu, Met
Ser, Thr,
Tyr, Trp
Asn, Gln
Asp, Glu
Lys, Arg, His

In additional embodiments the V_h region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and/or the V_l region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

In different embodiments focusing on an antibody, the antibody comprises, consists, or consists essentially of: (a) a heavy chain comprising a V_h region as described in this Section III, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄, and (b) a light chain comprising a V_l region as described above in this section III, and a human kappa C_L or human lambda C_L. In further embodiments: the antibody comprises, consists, or consists essentially of: (a) a heavy chain comprising a V_h region as described in this Section III, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁ or IgG₂ and (b) a light chain comprising a V_l region as described above in this Section III, and a human kappa C_L; and the heavy chain consists essentially of the amino acid sequence of SEQ ID NO: 22 and/or the light chain consists essentially of the amino acid sequence of SEQ ID NO: 23.

In additional embodiments the antigen-binding protein described herein has V_h and V_l regions providing an affinity K_D at least about 100 nM, preferably at least about 30 nM to the target antigen. Binding to the target antigen can be determined as described in Example 11, using an ORF0657n fragment from amino acids 42-486

Preferred binding proteins for the different embodiments are an antibody. More preferably the antibody is isolated or a monoclonal antibody.

IV. Protein Production

Antigen binding protein are preferably produced using recombinant nucleic acid techniques or through the use of a hybridoma. Recombinant nucleic acid techniques involve constructing

a nucleic acid template for protein synthesis. Hybridoma techniques involve using an immortalized cell line to produce the antigen binding protein. Suitable recombinant nucleic acid and hybridoma techniques are well known in the art. (See for example, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 2005, Harlow *et al.*, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988.)

Recombinant nucleic acid encoding an antigen binding protein can be expressed in a host cell that in effect serves as a factory for the encoded protein. The recombinant nucleic acid can provide a recombinant gene encoding the antigen binding protein that exists autonomously from a host cell genome or as part of the host cell genome.

A recombinant gene contains nucleic acid encoding a protein along with regulatory elements for protein expression. Generally, the regulatory elements that are present in a recombinant gene include a transcriptional promoter, a ribosome binding site, a terminator, and an optionally present operator. A preferred element for processing in eukaryotic cells is a polyadenylation signal. Antibody associated introns may also be present. Examples of expression cassettes for antibody or antibody fragment production are well known in art. (*E.g.*, Persic *et al.*, *Gene* 187:9-18, 1997, Boel *et al.*, *J. Immunol. Methods* 239:153-166, 2000, Liang *et al.*, *J. Immunol. Methods* 247:119-130, 2001, Tsurushita *et al.*, *Methods* 36:69-83, 2005.)

Due to the degeneracy of the genetic code, a large number of different encoding nucleic acid sequences can be used to code for a particular protein. The degeneracy of the genetic code arises because almost all amino acids are encoded by different combinations of nucleotide triplets or "codons". Amino acids are encoded by codons as follows:

A=Ala=Alanine: codons GCA, GCC, GCG, GCU

C=Cys=Cysteine: codons UGC, UGU

D=Asp=Aspartic acid: codons GAC, GAU

E=Glu=Glutamic acid: codons GAA, GAG

F=Phe=Phenylalanine: codons UUC, UUU

G=Gly=Glycine: codons GGA, GGC, GGG, GGU

H=His=Histidine: codons CAC, CAU

I=Ile=Isoleucine: codons AUA, AUC, AUU

K=Lys=Lysine: codons AAA, AAG

L=Leu=Leucine: codons UUA, UUG, CUA, CUC, CUG, CUU

M=Met=Methionine: codon AUG

N=Asn=Asparagine: codons AAC, AAU

P=Pro=Proline: codons CCA, CCC, CCG, CCU

Q=Gln=Glutamine: codons CAA, CAG

R=Arg=Arginine: codons AGA, AGG, CGA, CGC, CGG, CGU

S=Ser=Serine: codons AGC, AGU, UCA, UCC, UCG, UCU

T=Thr=Threonine: codons ACA, ACC, ACG, ACU

V=Val=Valine: codons GUA, GUC, GUG, GUU

W=Trp=Tryptophan: codon UGG

Y=Tyr=Tyrosine: codons UAC, UAU

Expression of a recombinant gene in a cell is facilitated using an expression vector. Preferably, the expression vector, in addition to a recombinant gene, also contains an origin of replication for autonomous replication in a host cell, a selectable marker, a limited number of useful restriction enzyme sites, and a potential for high copy number. Examples of expression vectors for antibody and antibody fragment production are well known in art. (*E.g.*, Persic *et al.*, *Gene* 187:9-18, 1997, Boel *et al.*, *J. Immunol. Methods* 239:153-166, 2000, Liang *et al.*, *J. Immunol. Methods* 247:119-130, 2001, Tsurushita *et al.*, *Methods* 36:69-83, 2005.)

If desired, nucleic acid encoding an antibody may be integrated into the host chromosome using techniques well known in the art. (*E.g.*, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 2005, Marks *et al.*, International Application Number WO 95/17516, International Publication Date June 29, 1995.)

A variety of different cell lines can be used for recombinant antigen binding protein expression, including those from prokaryotic organisms (*e.g.*, *E. coli*, *Bacillus sp.*, and *Streptomyces sp.* (or streptomycete) and from eukaryotic (*e.g.*, yeast, Baculovirus, and mammalian). (Breitling *et al.*, *Recombinant Antibodies*, John Wiley & Sons, Inc. and Spektrum Akademischer Verlag, 1999, Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004, Tsurushita *et al.*, *Methods* 36:69-83, 2005.)

Preferred hosts for recombinant antigen binding protein expression provide for mammalian post translational modifications. Post translational modifications chemical modification such as glycosylation and disulfide bond formation. Another type of post translational modification is signal peptide cleavage.

Proper glycosylation can be important for antibody function. (Yoo *et al.*, *Journal of Immunological Methods* 261:1-20, 2002, Li *et al.*, *Nature Biotechnology* 24(2):210-215, 2006.) Naturally occurring antibodies contain at least one N-linked carbohydrate attached to a heavy chain. (Yoo *et al.*, *Journal of Immunological Methods* 261:1-20, 2002.) Additional N-linked carbohydrates and O-linked carbohydrates may be present and may be important for antibody function. (*Id.*)

Different types of host cells can be used to provide for efficient post-translational modifications including mammalian host cells and non-mammalian cells. Examples of mammalian host cells include but are not limited to Chinese hamster ovary (Cho), HeLa, C6, PC12, Human Embryonic Kidney (HEK293) and myeloma cells. (Yoo *et al.*, *Journal of Immunological Methods* 261:1-20, 2002, Persic *et al.*, *Gene* 187:9-18, 1997.) Non-mammalian cells can be modified to replicate human glycosylation. (Li *et al.*, *Nature Biotechnology* 24(2):210-215, 2006.) Glycoengineered *Pichia pastoris* is an example of such a modified non-mammalian cell. (Li *et al.*, *Nature Biotechnology* 24(2):210-215, 2006.)

Preferred recombinant genes comprise a nucleotide sequence encoding an antibody variable region that binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region. A particular recombinant gene can encode for a protein containing one variable region or both a V_H and V_L region. The recombinant gene can also encode for antibody constant regions and hinge region. If desired, an antibody can be produced using a combination of recombinant genes, where one gene encodes for a light chain and the second gene encodes for a heavy chain.

Different embodiments are provided by the nucleic acid encoding a protein described in Section II or III *supra*. Examples of such embodiments are provided below.

In an embodiment directed to a V_H encoding region, the nucleotide sequence encodes a variable region comprising, consisting, or consisting essentially of: a first V_H CDR comprising, consisting, or consisting essentially of amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid; a second V_H CDR comprising, consisting, or consisting essentially of amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acid; and a third V_H CDR comprising, consisting, or consisting essentially of amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.

In an embodiment directed to a V_L encoding region, the nucleotide sequence encodes a variable region comprising, consisting, or consisting essentially of: a first V_L CDR comprising, consisting, or consisting essentially of amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid; a second V_L CDR comprising, consisting, or consisting essentially of amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and a third V_L CDR comprising, consisting, or consisting essentially of amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.

In additional embodiments: the V_H region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and the V_L region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

In different embodiments focusing on an antibody heavy and/or light chain, the recombinant gene encodes either or both a protein comprising, consisting, or consisting essentially of: (a) a heavy chain comprising a V_H region as provided in Section III *supra*, a human hinge, CH₁, CH₂, and CH₃ from an IgG1, IgG2, IgG3 or IgG4 subtype or (b) a light chain comprising a V_L region as provided in Section III *supra*, and a human kappa C_L or lambda C_L. In a further embodiment the heavy chain consists essentially of the amino acid sequence of SEQ ID NO: 22; and the light chain consists essentially of the amino acid sequence of SEQ ID NO: 23.

V. Applications of Antigen Binding Proteins.

Antigens containing certain ORF0657n regions can be used to provide a protective immune response against *S. aureus* infection. (Anderson *et al.*, International Publication No. WO

2005/009379, International Publication Date February 3, 2005.) An antigen binding protein recognizing an ORF0657n target region can be used to facilitate the production, characterization, or study of ORF0657n antigens and vaccines. Antigen binding protein recognizing appropriate epitopes can also have therapeutic applications.

Examples of different uses in the production, characterization, or study of ORF0657n related antigens and vaccines include:

- 1) Identifying the presence of an ORF0657n antigen, for example, by Western blot;
- 2) Identifying the presence of an ORF0657n antigen on a cell surface, for example, by flow cytometry. This is useful, for example, in determining expression on multiple strains of *S. aureus* as well as confirmation of knock-out mutants;
- 3) Passive protection experiments. The antibodies can be used in a lethal model to determine if a specific area of the ORF0657n protein confers protection;
- 4) An immunoassay. The assay can be used to monitor antigen quality, product production and stability;
- 5) As a control in mouse potency assays to monitor immunogenicity of an antigen vaccine product; and
- 6) Serology assays can utilize a monoclonal antibody in a competitive format to identify an immune response to ORF0657n derived antigen vaccinated patients.

Techniques for using antigen binding proteins, such as monoclonal antibodies, in the production, characterization, or study of a target protein are well known in the art. (See, for example, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 2005, Harlow *et al.*, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988, Harlow *et al.*, *Using Antibodies*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., Cold Spring Harbor Laboratory Press, 1999, Lipman *et al.*, *ILAR Journal* 46:258-268, 2005.)

In an embodiment of the present invention, the presence of an ORF0657n antigen in a solution, bound to a microsphere or on a cell is determined using an antigen binding protein. The ability of the binding protein to bind to a protein present in the solution or cell can be determined using different techniques such as a Western blot, enzyme-linked immunosorbent assay (ELISA), flow cytometry, and Luminex immunoassay.

VI. Treatment

Therapeutic and prophylactic treatment can be performed on a patient using an antigen binding protein binding to an appropriate target region. Therapeutic treatment is performed on those persons infected with *S. aureus*. Prophylactic treatment can be performed on the general population or a subset of the general population. A preferred subset of the general population are those persons at an increased risk of *S. aureus* infection.

A "patient" refers to a mammal capable of being infected with *S. aureus*. Preferably, the patient is a human. However, other types of mammals such as cows, pigs, sheep, goats, rabbits, horses, dogs, cats, monkeys, rats, and mice, can be infected with *S. aureus*. Treatment of non-human patients is useful in protecting pets and livestock, and in evaluating the efficacy of a particular treatment.

Persons with an increased risk of *S. aureus* infection include health care workers; hospital patients; patients with a weakened immune system; patients undergoing surgery; patients receiving foreign body implants, such as a catheter or a vascular device; patients facing therapy leading to a weakened immunity; and persons in professions having an increased risk of burn or wound injury. (*The Staphylococci in Human Disease*, Crossley and Archer (ed.), Churchill Livingstone Inc. 1997.)

In an embodiment, a patient is administered an antigen binding protein in conjunction with surgery or a foreign body implant. Reference to "surgery or a foreign body implant" includes surgery with or without providing a foreign implant, and providing a foreign implant with or without surgery. The timing of administration can be designed to achieve prophylactic treatment and/or therapeutic treatment. Administration is preferably started around the same time as surgery or implantation.

Guidelines for pharmaceutical administration in general are provided in, for example, *Remington's Pharmaceutical Sciences 20th Edition*, Ed. Gennaro, Mack Publishing, 2000; and *Modern Pharmaceutics 2nd Edition*, Eds. Banker and Rhodes, Marcel Dekker, Inc., 1990.

Pharmaceutically acceptable carriers facilitate storage or administration of an antigen binding protein. Substances used to stabilize protein solution formulations include carbohydrates, amino acids, and buffering salts. (Middaugh *et al.*, *Handbook of Experimental Pharmacology* 137:33-58, 1999.)

Antigen binding proteins can be administered by different routes such as intravenous, subcutaneous, intramuscular, or mucosal. Subcutaneous and intramuscular administration can be performed using, for example, needles or jet-injectors. Mucosal delivery, such as nasal delivery, can involve using enhancers or mucoadhesives to produce a longer retention time at adsorption sites. (Middaugh *et al.*, *Handbook of Experimental Pharmacology* 137:33-58, 1999.)

Suitable dosing regimens are preferably determined taking into account factors well known in the art including age, weight, sex and medical condition of the patient; the route of administration; the desired effect; and the particular compound employed. It is expected that an effective dose range should be about 0.1 mg/kg to 20 mg/kg, or 0.5 mg/kg to 5 mg/kg. The dosing frequency can vary depending upon the effectiveness and stability of the compound. Examples of dosing frequencies include biweekly, weekly, monthly and bimonthly.

VII. Examples

Examples are provided below further illustrating different features of the present invention. The examples also illustrate useful methodology for practicing the invention. These examples do not limit the claimed invention.

Example 1: Generation of Monoclonal Antibodies to ORF0657n

Monoclonal antibodies directed to ORF0657n (SEQ ID NO: 1) were generated using ORF0657n-C/e (SEQ ID NO: 2) or ORF0657n-H/y (SEQ ID NO: 3) as an antigen. The antibodies were identified and characterized by ELISA and flow cytometry.

Mice and Immunizations: Female BALB/c mice, 4-5 weeks old, were purchased from Taconic (Germantown, N. Y.). Mice were immunized intramuscularly (i.m.) on days 0, 7, and 21, with 20 µg of *E. coli* produced ORF0657n-C/e antigen or Yeast expressed ORF0657n-H/y antigen, formulated on aluminum hydroxyphosphate adjuvant. (Anderson *et al.*, International Publication No. WO 2005/009379, International Publication Date February 3, 2005.) A final intravenous injection (i.v.) of 20 µg of protein in phosphate buffered saline (PBS) was given to mice three days prior to the fusion. Mice were sacrificed and the spleens removed for cell fusion.

Mab Production: Lymphocytes prepared from spleens were fused with the mouse myeloma partner SP2/0-Ag14 (ATCC 1581) by polyethylene glycol 1500 (Boehringer Mannheim) at a ratio of 3:1. The fusions were plated into 96-well flat-bottomed microtiter plates in Dulbecco's Modification of Eagle's Medium, high glucose, pyruvate (DMEM) containing 20% fetal bovine serum, hypoxanthine (10^{-4} M), thymidine (10^{-5} M), Aminopterin (4×10^{-7} M) was added 24 hours later. Supernatants from growing hybridomas were screened by ELISA for reactivity to ORF0657n as described below. Positive wells were cloned by limiting dilution and retested for ELISA reactivity. Monoclonal antibodies were classified with an antibody-isotyping kit (Roche Diagnostics Corporation, Indianapolis, IN).

ELISA: Costar medium binding microtiter plates were coated overnight at 2-8°C with 50 nanograms per well of *E. coli* expressed SEQ ID NO: 2 in PBS. The plate was washed three times with PBS, 0.05% Tween20 and blocked with 1% Bovine serum albumin, PBS, 0.05% Tween20 (assay diluent) for at least 1 hour. The plate was washed as before and supernatants from the fusion wells or cloned hybridomas were added and allowed to incubate for 2 hours at room temperature. The plate was washed as before and a Goat anti-mouse IgG (H+L)-HRP conjugate (Zymed) (1:8000 in assay diluent) added and allowed to incubate for 1 hour at room temperature. Assay plates were developed with TMB substrate, the reaction stopped with 2.0 N H₂SO₄ and read in a plate reader at OD 450 nm. Wells were considered positive that had an optical density at 450 nm of >1.0.

Flow Cytometry: Prepared glycerol stocks of *S. aureus* passaged under iron-starved conditions (in RPMI) were used to evaluate mAb for ORF0657n binding. Frozen glycerol stock cells were thawed and resuspended in PBS; 1% bovine serum albumin; 0.1% sodium azide, 0.2% Pig IgG (Sigma) (PAAG) to a concentration of 5×10^7 CFU/50 µl. A 50 µl aliquot of the cells were placed in a 1.5 ml Eppendorf tube per reaction. Fifty microliters of the hybridoma culture were added to each reaction tube and incubated for 1 hour at room temperature. The cells were washed by adding 1 mL of phosphate buffered saline; 1% bovine serum albumin; 0.1% sodium azide (PAA) to the tube. The cells

were pelleted by centrifugation (5500 rpm, 5 minutes). The supernatant was removed and the cells were mixed with 100 μ L of secondary antibody (FITC-labeled goat anti-mouse Ig (BD Pharmingen) diluted 1:100 in PAAG). Incubation was for 1 hour at room temperature in the dark. After incubation, 1 mL PAA was added to the reaction mixture, the cells were pelleted (5500 rpm, 5 minutes) and supernatant removed. The pellets were resuspended in 1 mL of PBS and transferred to 12 x 75 mm tubes for FAC analysis.

Tubes were run on a BD-FACSCalibur flow cytometer instrument gated for bacterial cells and measuring the amount of FITC associated with the cells. A standard antibody with known binding to the surface of *S. aureus* was run in every assay. A negative control was run as cells and the secondary conjugate alone. Hybridoma wells were considered positive if the geometric mean value was greater than 30.

Two separate fusions resulted in a panel of twelve monoclonal antibodies (mAb). All of the mAbs were reactive in ELISA (Table 2). Ten of the twelve mAbs bound to the surface of bacteria as evidenced by flow cytometry. All of the mAbs were positive by Western Blot analysis with the wild type protein.

Table 2

mAbs/cell lines Fusion #1	mAbs/cell lines Fusion #2
1) 2H2.B8 IgG1	
2) 8H6.E11.H3 IgG2a*	
3) 7H2.C11 IgG1*	
	4) 2E12.A8 IgG1
	5) 8A8.B4 IgG1
	6) 3G11.D5 IgG1
	7) 13G11.C11 IgG1
	8) 13C7.D12 IgG1
	9) 1G3.B3 IgG1
	10) 9H3.E4 IgG1
	11) 3B7.G8 IgG1
	12) 3G12.A4 IgG1

* Not reactive in flow cytometry. Fusion #1 was generated from *E. coli* produced ORF0657n-C/e antigen. Fusion #2 was generated with Yeast expressed ORF0657n-H/y antigen.

Example 2: Class Switching mAbs

All of the mAbs isolated that bound to the native antigen were of the IgG1 isotype. These antibodies were class switched to an IgG2b isotype by selecting for shift variants (Spira *et al*, *J. of Immunological Methods*, 74:307-315, 1985). A suitable immunoassay was developed using an IgG2b conjugate and the cell line was plated at a high density. Somatic cell mutations were selected, enriched

and then cloned. The binding site of the switched mAb remained identical to the original mAb, but switching to an IgG2b subtype gave a more favorable isotype (initiating the complement cascade) in the passive protection studies.

Table 3 Class Switched mAbs

IgG1 isotype	IgG2b isotype
2H2.B8	2H2.BE11
2E12.A8	2E12.BG1
8A8.B4	8A8.BF9
3G11.D5	3G11.BE5
13G11.C11	13G11.BF3
13C7.D12	13C7.BC1
1G3.B3	1G3.BD4
9H3.E4	9H3.BE4

Example 3: Binding Inhibition Studies with Native Antigen

Purified antibodies were labeled with Alexafluor-488 using a mAb labeling kit (Molecular Probes) according to the manufacturer's instructions. The amount of mAb that would just saturate the surface of RPMI-grown bacterial cells was determined for both the labeled and unlabeled mAbs. Each of the mAbs in Table 3 (1st column) were used labeled and unlabeled.

The inhibition assay was performed by first incubating 5×10^7 cells with the unlabeled mAb at a concentration that would saturate the surface of the cells. This reaction was incubated at room temperature for 1 hour. After this incubation, the reactions were washed with 1 ml of PAA and spun at 6,000 RPM for 5 minutes in a microcentrifuge (Hermle). The supernatant was removed down to ~50 ul and the cells were resuspended in 100 ul of PAAG containing the amount of directly labeled mAb that would just saturate the surface of the cells. After this incubation, the reactions were washed with 1 ml of PAA and spun at 6,000 RPM for 5 minutes in a microcentrifuge (Hermle). The supernatant was removed down to ~50 ul and the cells were resuspended in 1 ml of PBS and transferred to 12 x 75 mm tubes for FAC analysis. As controls, separate reactions with the unlabeled mAb were measured with a secondary Alexafluor-488 conjugated goat anti-mouse IgG (H+L) (Molecular probes, 1:400 in PAAG) to determine that this mAb was bound to the surface. A positive control was also performed that had only the labeled mAb with the cells. If the unlabeled mAb bound to the same epitope as the labeled mAb then there would be no or low fluorescent reactivity associated with the cells. If the unlabeled mAb bound to a different epitope than the labeled mAb then the level of reactivity associated with the surface would be equivalent to the labeled mAb only control cells.

The panel of monoclonal antibodies fell into four reactive groups by inhibition studies:

Table 4

Group I	Group II	Group III	Group IV
2H2.B8	9H3.E4	13G11.C11	2E12.A8
8A8.B4	1G3.B3		13C7.D12
	3G11.D5		

Example 4: Binding Studies with Denatured Antigen and Altered Antigens

ORF0657n altered proteins were used to further characterize binding. Nucleic acid encoding ORF0657n was initially cloned into the expression vector pET-28a (Novagen) and expressed in *E. coli* with a C-terminal 6X his tag (SEQ ID NO: 2). The expression vector with the cloned gene was subjected to mutagenesis using Stratagene's QuikChange XL Site-Directed Mutagenesis Kit following the manufacturer's instructions. The gene was mutated with specific sequential amino acid changes. The resulting plasmid was transformed into Stratagene's XL10-Gold competent cells following the manufacturer's protocol. Plasmids were isolated from transformants using Qiagen's QIAprep Spin Miniprep Kit. Transformants were screened by sequencing using ABI's 310 DNA Sequencer. Plasmid from the transformant exhibiting the greatest number of base changes was transformed into the expression host HMS174(DE3) (Novagen). Transformants were expressed following Novagen's instructions.

Different ORF0657n altered proteins were used to determine the diversity of the ORF0657n mAbs (SEQ IDs 4-19). These proteins were screened with the 10 different mAbs in dot blots using standard procedures. Positive/negatives were confirmed by Western blots using standard procedures. By this approach antibodies were grouped according to their binding profile. Seven of the antibodies resolved to three groups; the three remaining antibodies (2H2.B8, 8A8.E11.H3 and 13G11.C11) had profiles that were similar but not identical to each other (Table 5).

TABLE 5: Binding of ORF0657n specific mAbs to ORF0657n mutant proteins detected by Western blot

	Group III		Group II			Group IV		Group I		
SEQ ID NO:	3G11.C11	3G12.A4	3B7.G8	1G3.B3	9H3.E4	2E12.A8	13C7.D12	2H2.B8	8A8.E11.H3	13G11.C11
1	+	+	+	+	+	+	+	+	+	+
2	+	+	+	+	+	+	+	+	+	+
3	+	+	+	+	+	+	+	+	+	+

TABLE 5: Binding of ORF0657n specific mAbs to ORF0657n mutant proteins detected by Western blot

	Group III		Group II			Group IV		Group I		
SEQ ID NO:	3G11.C11	3G12.A4	3B7.G8	1G3.B3	9H3.E4	2E12.A8	13C7.D12	2H2.B8	8A8.E11.H3	13G11.C11
4	+	+	+	+	+	+	+	+	+	+
5	+	+	+	+	+	+	+	+	+	+
6	+	+	+	+	+	-	-	+	+	+
7	+	+	+	+	+	-	-	+	+	+
8	+	+	+	+	+	-	-	+	+	+
9	+	+	+	+	+	-	-	-	+	+
10	+	+	+	+	+	-	-	-	+	+
11	-	-	W	W	W	-	-	-	-	W
12	-	-	W	W	W	-	-	-	-	W
13	-	-	-	-	-	-	-	-	-	-
14	+	+	+	+	+	+	+	+	+	+
15	+	+	+	+	+	+	+	+	+	+
16	+	+	+	+	+	+	+	-	+	+
17	+	+	+	+	+	+	+	-	+	+
18	+	+	+	+	+	+	+	-	+	-
19	+	+	+	+	+	W	W	-	-	+

+, Antibody bound to protein in a Western; -, Antibody did not bind to protein by Western; W, Weak binding of antibody to protein detected by Western. Antibodies were grouped according to hybridization profile. A dotted line is used where similar, but not identical profiles were obtained.

Example 5: BIAcore Studies

In BIAcore studies the mAbs were examined by "footprint analysis" using purified ORF0657n-H/y as the antigen. Pair-wise binding experiments were conducted using real-time biomolecular interaction analysis via BIACORE[®]. BIACORE[®] incorporates microfluidics technology and surface plasmon resonance (SPR) to detect changes in mass by monitoring changes in the refractive index of a polarized light aimed directly at the surface of a carboxyl methyl dextran coated (CM5) sensor chip. The changes in response, measured in Response Units, can be correlated to the amount of bound analyte (*i.e.* antigen or antibody).

An anti-staphylococcal antibody (mAb 13C7.D12) was covalently bound (immobilized) on the surface of the CM5 sensor chip. The immobilized Ab was exposed first to the ORF0657n protein and subsequently to a pair of antibodies in a matrix format. After each cycle of ORF0657n protein + antibody pair, the surface of the sensor chip was regenerated back to the immobilized mAb 13C7.D12

using 20 mM HCl. Eight antibodies were tested against the ORF0657n protein in a matrix format so that all combinations of each antibody pair could be analyzed. The matrix design for mAb pairs used in this experiment is summarized in Table 6.

Table 6. Summary of Antibodies Tested in 8x8 Matrix

Cycle #	First Antibody	Second Antibody			
		Flow Cell 1	Flow Cell 2	Flow Cell 3	Flow Cell 4
1	N/A Immobilization	13C7.D12	13C7.D12	13C7.D12	13C7.D12
2	2H2.B8	2H2.B8	13C7.D12	8A8.B4	9H3.E4
3	2H2.B8	13G11.C11	2E12.A8	1G3.B3	3G11.D5
4	13C7.D12	2H2.B8	13C7.D12	8A8.B4	9H3.E4
5	13C7.D12	13G11.C11	2E12.A8	1G3.B3	3G11.D5
6	8A8.B4	2H2.B8	13C7.D12	8A8.B4	9H3.E4
7	8A8.B4	13G11.C11	2E12.A8	1G3.B3	3G11.D5
8	9H3.E4	2H2.B8	13C7.D12	8A8.B4	9H3.E4
9	9H3.E4	13G11.C11	2E12.A8	1G3.B3	3G11.D5
10	13G11.C11	2H2.B8	13C7.D12	8A8.B4	9H3.E4
11	13G11.C11	13G11.C11	2E12.A8	1G3.B3	3G11.D5
12	2E12.A8	2H2.B8	13C7.D12	8A8.B4	9H3.E4
13	2E12.A8	13G11.C11	2E12.A8	1G3.B3	3G11.D5
14	1G3.B3	2H2.B8	13C7.D12	8A8.B4	9H3.E4
15	1G3.B3	13G11.C11	2E12.A8	1G3.B3	3G11.D5
16	3G11.D5	2H2.B8	13C7.D12	8A8.B4	9H3.E4
17	3G11.D5	13G11.C11	2E12.A8	1G3.B3	3G11.D5

To normalize for the amount of antigen initially bound (captured) in each run, the following ratio for each test antibody/antigen complex is calculated:

$$= \frac{\text{Test Antibody Response Units} * 1000}{\text{ORF0657n protein Response Units}} \quad \text{or} \quad \frac{\text{mRU}_{\text{Ab}}}{\text{RU}_{\text{Ag}}}$$

The percentage of available epitope remaining for each antibody can be calculated for the mapping pair as follows:

$$= \frac{(\text{mRU}_{\text{Ab}} (\text{when } 2^{\text{nd}} \text{ Ab}) / \text{RU}_{\text{Ag}}) * 100}{(\text{mRU}_{\text{Ab}} (\text{when } 1^{\text{st}} \text{ Ab}) / \text{RU}_{\text{Ag}})} \quad \text{or} \quad \begin{array}{l} \% \text{ Remaining} \\ \text{(calculated for each Ab)} \end{array}$$

Figure 2 illustrates matrix resulting outlining the reactivities of the monoclonal antibodies in a pair-wise binding study. The panel of monoclonal antibodies fell into three reactive areas by the BIAcore® method (See Table 7).

Table 7

Group I	Group II	Group III
2H2.B8	13G11.C11	13C7.D12
8A8.B4	3G11.D5	2E12.A8
9H3.E4		
1G3.B3		

Example 6: Protection Studies with Passive Immunization in a Murine Sepsis Model

The monoclonal antibodies mAb 2H2.BE11 and mAb 13C7.BC1 were tested for their ability to provide protection against *S. aureus* infection. These antibodies recognize different epitopes on the ORF0657n protein. Controls included an isotype matched mAb and PBS-only.

The mAbs or PBS were administered intraperitoneally (i.p.) 20 hours prior to bacterial challenge. Mice were then challenged with a LD₈₀₋₉₀ dose of *S. aureus* Becker i.v. and monitored for survival. Each experiment was repeated three times with groups of 10 or 20 mice and was monitored for 10 days. The half life for the monoclonal antibodies in uninfected BALB/c mice is approximately eight days. A dose of 0.5 mg was found to be optimal. The results of experiments with the two monoclonal antibodies are presented in Figures 3A-C, 4A, 4B, and 5A-C.

Whereas the mAb 13C7.BC1 significantly improved survival at day 10 compared to the controls in one experiment, in the other 2 repetitions the overall survival rate was similar to that of the controls (Figures 3A-3C). However, compared to controls, there was delay in the time to death of the mAb 13C7.BC1 treated mice within this 10 day period. A similar trend in delay of time to death of the mAb 2H2.BE11 treated mice was also noted in two of the three experiments (Figures 5A-5C).

The effect of mAb 13C7.BC1 was also examined using a recent *S. aureus* clinical isolate UK58 (Figures 4A and 4B). This strain was minimally passaged from an abscess site in a patient. In two independent experiments, the results show a delay in time to death with the UK58 challenge.

Antibody persistence studies cannot be evaluated in the LD₈₀₋₉₀ model due to the rapid rate of death. Therefore, a sub-lethal challenge model was run. In the sub-lethal model the challenge dose used is 10% of that used for the LD₈₀₋₉₀ model. The sub-lethal challenge model was monitored over a four day period. Groups of 22 mice received 0.5 mg doses of either mAb 13C7.BC1 or isotype control mAb (6G6) 20 hours prior to i.v. bacterial challenge with 5×10^7 CFU of *S. aureus* Becker. Two animals from each group were sacrificed just prior to challenge (T=0) to determine the mAb levels in the

serum at the time of challenge. At 2, 24, 48, 72 and 96 hours post challenge, four mice from each group were sacrificed and serum mAb levels determined.

From this sub-lethal challenge experiment, the half life of mAb 13C7.BC1 in *S. aureus*-infected mice was estimated to be approximately one-day. In contrast, the half life of the isotype control mAb was estimated to be greater than four days (data not shown). These data point to a specific reduction of mAb 13C7.BC1 in *S. aureus* challenged mice, which appears to be exhausted well before the ten day period monitored in the lethal model.

In six of the eight experiments illustrated in Figures 3A-C, 4A, 4B, and 5A-C, improved survival was observed through approximately three days for the groups receiving the mAb administration. These results provide an indication that such mAbs have a positive effect on the survival rate of *S. aureus* challenged mice.

Example 7: Protection Studies with Passive Immunization in a Murine Indwelling Catheter Model

A murine indwelling catheter model was used with mAb 2H2.BE11. The *S. aureus* strain used in this model was the clinical isolate MCL8538. This strain was selected as lower inocula could be administered while still getting reproducible colonization of catheters compared to *S. aureus* Becker, the strain used in the murine sepsis model.

ICR mice had catheters (PE50 silicone rubber) surgically implanted into the jugular vein, held in place with sutures, and exiting with a port on the dorsal midline of the mouse. Mice were rested 9-11 days post surgery. At 24 hours prior to challenge, mice were passively immunized with a single injection of 600 mcg of murine monoclonal antibody 2H2.BE11 administered i.p. At day 0, mice were challenged with *S. aureus* MCL8538 administered i.v. The inoculum dose was 2×10^5 CFU in 100 μ l volume (Experiments 1 to 3). This low dose was found to clear spontaneously from the catheters after 4 days. Therefore, catheters were assessed for bacteria at 24 hours post challenge. At that time, mice were sacrificed and catheters harvested. The presence of bacteria on the catheters was assessed by culturing the entire catheter on TSA. If any sign of outgrowth was observed on the plate the catheter was scored as culture positive.

In two of the first three experiments, the number of culture negative catheters was significantly lower in mice passively immunized with antibody 2H2.BE11, as compared to the isotype control antibody. A fourth experiment was performed using a larger inoculum dose. In this more rigorous challenge, the dose was determined to be one in which 100% of catheters were reproducibly infected, and this infection was not spontaneously cleared by control mice (monitored over 7 days). In experiment 4, with the larger inoculum size, again, significantly fewer catheters were found to be infected in mice injected with antibody to 2H2.BE11, compared with the isotype control. Results of the four experiments are summarized in Table 8.

Table 8: Number Of Culture Negative Catheters Obtained In 4 Independent Passive Transfer Experiments Using a Murine Indwelling Catheter Model

Monoclonal	Number of Culture-Negative Catheters					p-value
	Exp#1	Exp#2	Exp#3	Exp#4	Total	
2H2.BE11	3 of 4 (75%)	6 of 8 (75%)	4 of 10 (40%)	4 of 9 (44%)	17/31 (54%)	0.0187
Isotype matched control	1 of 4 (25%)	3 of 8 (38%)	4 of 10 (40%)	0 of 9 (0%)	8/31 (25%)	

Groups of ICR mice with indwelling catheters were injected i.p. with 600 mcg of murine monoclonal antibody 24 hours prior to challenge, all monoclonals of the IgG2b isotype

Example 8: Ex-Vivo Pre-Opsonization of Bacteria Using anti-ORF0657n Monoclonal Antibodies 2H2.B8 (IgG1), 2H2.BE11 (IgG2b), or 13C7.IgG2b or Isotype Matched Control mAbs

To test whether monoclonal antibodies to ORF0657n are opsonic, passive protection experiments were conducted in which a lethal dose of *S. aureus* was pre-opsonized with the monoclonal antibodies 2H2.B8, 2H2.BE11, or 13C7.IgG2b, or an isotype matched control monoclonal antibody. Pre-opsonized bacteria were then administered to mice i.p. Bacteria used in these experiments were *S. aureus* RN4220 (wild type) or RN4220.0657n. The RN4220.0657n bacteria were engineered to express ORF0657n in the absence of control by the FUR box. Therefore, they could be grown in the presence of iron and still express ORF0657n antigen on their surface. Alternatively, RN4220 (wild type) was passed 2X in a low iron medium RPMI to induce expression of 0657n on the bacteria surface.

A quantity of bacteria sufficient for 6 Balb/c mice (6 X LD₁₀₀) was incubated with 800 µg IgG at 4 °C for 1 hour, with gentle rocking. Bacteria were then pelleted and any unbound mAb removed. Antibody-opsonized bacteria were re-suspended in 2.4 mL of PBS, and 0.4 mL (1 X LD₁₀₀) was injected into each of five mice. After challenge, each inoculum was quantitated by plating on TSA to insure that equivalent CFU was given to all groups of mice and that the mAbs had not aggregated the bacteria. Survival was monitored for 3 days post challenge. Since the target antigen must be present on the surface of the bacteria for this procedure to be effective, care was taken to ensure that 0657n was expressed on the bacteria prior to opsonization. ORF0657n expression was monitored by flow cytometry using mAb 2H2.B8. The dose of opsonized bacteria injected into each mouse was 2-4 X 10⁹ CFU RN4220.0657n/mouse, or 1-2 X 10⁹ CFU RN4220(2X RPMI)/mouse.

When pre-opsonized with either 2H2.B8 or 2H2.BE11, but not an isotype matched control mAb, mice were protected from death from a lethal dose of RN4220.0657n staphylococci. The experiment was repeated twice for the IgG1 isotype and three times for the IgG2b isotype with similar results (Table 9A).

Table 9A: Ex-vivo Protection with Anti-0657n mAb

Monoclonal	Exp 1 Surviving Mice	Exp 2 Surviving Mice	Exp 3 Surviving Mice	Total
2H2.BE11 (IgG2b)	5	4	5	93% (14/15)
6G6.A8 (IgG2b)	1	0	1	13% (2/15)
PBS	1	2	0	20% (3/15)
2H2.BE11 (IgG1)	ND	4	5	90% (9/10)
10B4.H4 (IgG1)	ND	1	1	20% (2/10)

Five mice were used in each experiment. Challenge strain RN4220.0657n.pYZ119. Dose: $2-4 \times 10^9$ CFU. Test mAbs: murine anti-0657n 2H2.BE11 (IgG2b); 2H2.B8 (IgG1).

When pre-opsionized with either mAb 2H2.B8 but not an isotype matched control mAb, mice were protected from death from a lethal dose of RN4220 (2X RPMI) staphylococci. The experiment was repeated six times with similar results (Table 9B).

Table 9B: Ex-vivo Protection with Anti-0657n mAb

Monoclonal	# Tests	Aggregate	% Survival
2H2.B8	6	30/30	100%
10B4.IgG1 Isotype control	6	2/30	7%
13C7.IgG2b	2	0/10	0%
6G6.IgG2b Isotype control	2	0/10	0%

Murine anti-0657n 2H2 was very effective in preventing death in this lethal model. The 13C7 mAb was not effective in this model (as opposed to the previously described model illustrated in Figures 3-6). All (2H2.BE11, 2H2.B8 and 13C7.IgG2b) of the anti-0657n mAb's bind RN4220 (as demonstrated using flow cytometry) and all have opsonizing activity in the *in vitro* OPA assay. This model reflects an additional requirement for epitope specificity for enhancing survival in the peritoneum of the mouse.

Example 8: Epitope mapping studies performed with 2H2 mAb

The experiments described in this example provide evidence that the monoclonal antibody 2H2.BE11 recognizes a conformational epitope within ORFO657n. The experiments localized the minimal sequence within ORFO657n required for displaying the conformational epitope in a three dimensional structure recognized by 2H2 mAb. In addition, the experiments identified distinct lysine

residues within the minimal sequence of ORFO657n that become protected from reacting with small molecules when 2H2 mAb is bound to ORFO657n.

The potential ability of 2H2 mAb to recognize linear epitopes of typically 9 to 14 amino acids in length within the sequence of ORFO657n was investigated using epitope extraction and starting with an ORFO657n fragment from amino acid 42 to amino acid 486 of SEQ ID NO: 1 ("ORFO657t"). In detail: 30 ug of 2H2 mAb were immobilized by chemical cross linking to 10 mg of cyanogen bromide activated sepharose (Amersham cat. No. 17 0430 01) for each of the epitope extraction experiments. Proteolytic digests of the ORFO657t were generated with GluC (Roche Applied Science cat. No. 11 420 3997 001), Asp-N (Roche Applied Science cat. No. 11 054 589 001) or Chymotrypsin (Roche Applied Science cat. No. 11 418 467 001) and characterized by 1D/LC-MS/MS on a linear ion trap (LTQ – Thermo Electron Inc). In three individual experiments 8.4 µg of the characterized proteolytic digest from any protease was allowed to react with the immobilized antibody. Unbound peptides were washed off the antibody cross-linked beads. Potentially bound peptides were eluted with low pH and characterized by 1D/LC-MS/MS. None of the generated proteolytic peptides were recognized with high efficiency and specificity by 2H2 mAb, providing a strong indication that 2H2 mAb did not recognize a linear epitope.

The finding that 2H2 mAb did not recognize a linear sequence of ORFO657n was corroborated by a limited chemical cleavage experiment. ORFO657t was chemically cleft with CNBr for 2 hours. The resulting cleavage products were analyzed by SDS-PAGE. SDS-PAGE analysis showed 5 major bands with molecular weights of approximately 42 kDa, 35 kDa, 25 kDa, 15 kDa and 10 kDa. A Western Blot analysis with 2H2 mAb clearly showed that only the 42 kDa band was recognized by 2H2. All bands were excised from the SDS-PAGE, in-gel digest was performed, and the resulting peptides that were identified by tandem mass spectrometry were matched to corresponding sequences in ORFO657t. The result of the analysis of the major bands is shown in Table 10:

Table 10

CNBr cleavage	Binds to 2H2 mAb	ORFO657t	Calculated MW kDa
Band 42 kDa	yes	[001-356]	40.7
Band 35 kDa	no	[001-323]	36.7
Band 25 kDa	no	[001-214]	23.9
		[116-302]	21.9
Band 15 kDa	no	[215-356]	16.8
		[303-446]	16.6
Band 10 kDa	no	[114-214]	11.7
		[215-302]	10.39
		[357-446]	10.28

The importance of a fragment with a molecular weight of ~ 42 kDa was confirmed by epitope excision. In detail, 210 µg of 2H2 mAb was immobilized by chemical cross linking to 50 mg of cyanogen bromide activated sepharose (Amersham cat. No. 17 0430 01) for each of the epitope excision experiments. Then, 50 µg of intact ORF0657t was allowed to bind to the immobilized antibody and non-bound ORF0657t washed off by intensive washing with phosphate buffered saline. In three independent experiments proteases Glu-C, Trypsin and a sequential combination of Glu-C, AspN, Trypsin, Chymotrypsin, and Carboxy-peptidase Y were added for 5 hours or one hour per protease in the sequential combination. Peptides that were excised by the proteases during the incubation were thoroughly washed away and ORF0657t fragments that specifically bound to 2H2 mAb released with SDS loading buffer.

Fragments that specifically bound to 2H2 mAb were analyzed by SDS-page. All three of the epitope excision experiments showed exclusively one band with a molecular weight between 40 and 42 kDa in the SDS-Page analysis. Bands binding to 2H2 mAb were confirmed by Western Blot analysis. The epitope excision experiment was repeated for the Glu-C protease. This time the fragment of ORF0657t that specifically bound to 2H2 mAb was released with acidic conditions and analyzed by 1D/LC-MS/MS on a linear ion trap (LTQ, Thermo Electron). The eluted sample showed a signal (total ion count) with the expected intensity at 82-87 minutes (40%– 45% acetonitrile) and multiple charge states ($[M+67\text{ H}]^{67+}$ to $[M+30\text{ H}]^{30+}$) that deconvoluted to 42.628 kDa. A possible fragment of ORF0657t corresponding to this particular mass is sequence [012-382] of ORF0657t with a molecular weight of 42.6 kDa.

To determine which lysine residues of ORF0657t are protected from chemical reactions upon binding of 2H2 mAb, chemical labeling experiments were performed with sulfo-NHS-acetate (Pierce Cat. No. 26777) using three different experimental conditions in the presence or absence of 2H2 mAb. See Table 11.

Table 11

Experiment	1	2	3
molar excess 2H2 mAb	0 or 3	0 or 3	0 or 3
molar excess sulfo-NHS acetate	25	500	75
Reaction temperature °C	room temperature	15	37
Reaction time	1 hour	30 minutes	2 hours

For each experiment, reaction products produced with 0 or 3 molar excess 2H2 mAb were incubated with one of three proteases resulting in 2 x 9 reaction mixtures. Experiment 1 employed

GluC, AspN and Trypsin. Experiments 2 and 3 employed GluC, AspN, and Chymotrypsin. The proteolytic peptides were then analyzed by 1D/LC-MS/MS. For each of the reactions a ratio of acetylated and non-acetylated lysine residues was calculated based on the area under curve of the total ion count (TIC) of the individual peptides. Obtained ratios were then compared between the pairs (with and without 2H2 mAb) for identical reaction conditions. A global analysis was performed for all three reaction conditions to identify lysine residues within ORF0657t that are maximally shielded upon binding of ORF0657t to 2H2 mAb. The chemical labeling experiment described above identified K76, K257 and potentially K443 as being most protected upon binding of 2H2 mAb. Protection against chemical labeling is likely due to direct binding. However, it is possible that such protection could be due to binding in close proximity to the protected sites or by long range structural changes within ORF0657nI

In summary, the above described experiments provide clear evidence that the epitope within ORF0657t that is recognized by the 2H2 mAb is conformational. The fragment of ORF0657t that is recognized by 2H2 mAb has an N-terminus located between amino acids 1 and 115 of ORF0657t and a carboxyl terminus located between amino acids 323-357 of ORF0657t. Even though it can not be excluded that protection from chemical labeling upon binding of 2H2 mAb is influenced by long range structural changes, it is very likely that areas in close proximity to Lysine 76 and Lysine 275 participate in direct antibody interaction.

Example 9: 2H2 mAb Sequence Identification

Identification of the variable light (V_L) and variable heavy (V_H) sequences of hybridoma expressed 2H2 IgG was accomplished by combining the degenerative primer PCR /overlap extension cloning process for single chain variable fragments (scFv) assembly (Krebber *et al. JIM 201(1):35-55*, 1997), with high throughput screening of soluble scFv fused to a human kappa light chain constant domain or scAb material via Biacore. This allowed for fine discrimination of mutations in V_L frameworks 1, 4 and V_H frameworks 1, 4 generated by the degenerative primer method.

Briefly, RNA material was purified from the hybridoma cell line using standard methods from a Total RNA KitTM (Ambion Inc.). This material was then reverse transcribed to cDNA and utilized as template in PCR to amplify the variable regions. The conditions for the PCR amplification of the V_L and V_H chains was based upon the protocol described by Krebber *et al. JIM 201(1):35-55*, 1997. The primers are designed such that a (Gly4Ser)₄ linker (SEQ ID NO: 32) is added which provides domains for a third PCR reaction in which the V_H and V_L are overlapped to create a V_L -(Gly4Ser)₄- V_H scFv.

The first set of PCR reactions to amplify the variable chains individually, were carried out in a volume of 100 μ l containing 5 μ l of the cDNA reaction, 2 μ M each of the forward and reverse primer sets for amplification of V_L and V_H , and a high fidelity PCR master mix. The reactions were denatured for 4 minutes at 94°C followed by 30 cycles of 30 seconds at 94°C, 30 seconds at 50°C, 1

minute at 72°C, and finished at a final cycle of 5 minutes at 72°C. The full length PCR products were gel purified.

To construct the full length product a third PCR reaction was done to assemble to scFv from the amplified V_H and V_L material. In a volume of 100 µl approximately 20 ng each of V_H and V_L DNA and a high fidelity PCR master mix was denatured for 5 minutes at 94°C, followed by 3 cycles of 30 seconds at 94°C, 30 s at 60°C, and 30 seconds at 72°C in the absence of primers. The modified PCR primers, SEQ ID NO: 33 and SEQ ID NO: 34 were added at a final concentration of 1 µM, and 30 cycles of 30 seconds at 94°C, 1 minutes at 60°C, and 1 minute at 72°C were performed, followed by 7 minutes at 72°C. The expected full length scFv PCR products were gel purified.

The amplified scFv material was cloned into the MP16 soluble expression vector for scAb production (Hayhurst *et al.*, *JIM* 276(1-2):185-196, 2003) and sequence analysis. A single restriction enzyme digest with SfiI was used for directional cloning into the MP16 vector. Clones with apparent full length variable heavy and variable light chains present were then expressed as scAbs in XL1-Blue cells and recovered from the periplasm using a standard osmotic shock procedure. Briefly, clones were grown at 37°C overnight in growth media containing 2% glucose and 100 µg/ml ampicillin in a 96 well format. 20 µl of the overnight culture was transferred to new media containing 0.1% glucose and 100 µg/ml ampicillin and grown until an OD₆₀₀ of 0.6 was reached. The cells were induced for scAb expression by adding IPTG at a final concentration of 0.5 mM and incubated overnight while shaking at 150 rpm, at room temperature. The scAbs were purified from the cells using a Qiagen Ni-NTA superflow robotic procedure.

To analyze each scAb periplasmic preparation for binding activity to ORF0657t, a Biacore3000 surface plasmon resonance (SPR) instrument (Uppsala, Sweden) was utilized. Standard EDC/NHS coupling was used to covalently mobilize approximately 250 resonance units of the 0657t antigen directly to the experimental flow cell surface of a CM5 sensor chip. A reference flow cell surface was activated and deactivated without coupling of protein. Each preparation was then run over the surface and association and dissociation of the scAb to antigen was measured. The surfaces were regenerated between runs by a single injection of 10 mM HCl for 20 seconds at a flow rate of 20 µl/min, followed by a 2 minute stabilization period. All samples were run in duplicate and buffer only runs were used as controls. After screening 95 clones, a clone was selected based on its binding activity. The final 2H2 clone chosen was based upon its similar affinity for ORF0657t as the original hybridoma prepared IgG material as well as comparative sequence analysis.

The amino sequence of the 2H2 V_H (SEQ ID NO: 20) and V_L (SEQ ID NO: 21) were as follows:

2H2 V_H Amino Acid Sequence (SEQ ID NO: 20)

```

1   DVHLVESGPG LVAPSQNLSI TCTVSGFSLS RYGVHWVRQP PGKGLEWLGL
51  IWAGGVTIYN STLMSRLSIS KDSSKSQVFL KMNSLQIDDT AIYYCAREAS
101 RDHYFDYWGQ GTTLTVSS

```

2H2 V_L Amino Acid Sequence (SEQ ID NO: 21)

```

1   DIVMTQSPAI MSASPGKIT MTCSASSSVS YIYWYQQKSG TSPKRWIYDT
51  SKLASGVPFR FSGGGSGTSF SLTISSMEAE DAATYYCQOW SSNPLTFGAG
101 TKLEIK

```

The underlined portions are the CDR's. CDR's were identified based on the Kabat definition. The encoding nucleic acid sequence is provided by SEQ ID NO: 24 (V_H) and SEQ ID NO: 25 (V_L).

Example 10: 2H2 IgG Chimera Expression

The variable regions for 2H2 mAb were cloned from mouse hybridoma as described in Example 9. The sequences for the variable regions were PCR amplified and DNA encoding the heavy chain variable regions were fused in-frame with DNA encoding the IgG1 constant region whereas DNA encoding the light chain variable region were fused in-frame with DNA encoding the kappa constant region. The cloning procedure for the resulting antibody expression vectors is described below.

The variable regions were PCR amplified. PCR reactions were carried out in a volume of 25 µl containing high fidelity PCR master mix, template volume 1 µl and forward and reverse primers: 1 µl each. PCR condition was 1 cycle of 94°C, 2 minutes, 25 cycles of 94°C, 1.5 minutes; 60°C, 1.5 minutes; 72°C, 1.5 minutes and 72°C, 7 minutes; 4°C until removed and cloned in-frame with leader sequence at the 5'-end and constant region at the 3'-end using In-Fusion strategy. The following primers were used: Light chain forward, 5'- ACAGATGCCAGATGCGATATTGTGATGACCCAGTCT (SEQ ID NO: 28); Light chain reverse, 5'- TGCAGCCACCGTACGTTTATTTCCAGCTTGGTCCC (SEQ ID NO: 29); Heavy chain forward, 5'- ACAGGTGTCCACTCGGATGTGCACCTGGTGGAGTCA (SEQ ID NO: 30); and Heavy chain reverse, 5'- GCCCTTGGTGGATGCCGAGGAGACTGTGAGAGTGGT (SEQ ID NO: 31). The DNA sequences for all the clones were confirmed by sequencing.

The amino acid sequences deduced from DNA sequences are:

Mouse 2H2 Variable and Human Kappa Constant Region Amino Acid Sequence (SEQ ID NO: 22)

```

1   DIVMTQSPAI MSASPGKIT MTCSASSSVS YIYWYQQKSG TSPKRWIYDT
51  SKLASGVPFR FSGGGSGTSF SLTISSMEAE DAATYYCQOW SSNPLTFGAG
101 TKLEIKRTVA APSVFIFPPS DEQLKSGTAS VVCLLNNFYP REAKVQWKVD
151 NALQSGNSQE SVTEQDSKDS TYSLSSTLTL SKADYEKHKV YACEVTHQGL
201 SSPVTKSFNR GEC

```

Mouse 2H2 Variable and Human IgG1 Constant Region Amino Acid Sequence (SEQ ID NO: 23)

```

1   DVHLVESGPG LVAPSQNLSI TCTVSGFSLS RYGVHWVRQP PGKGLEWLGL
51  IWAGGVTIYN STLMSRLSIS KDSSKSQVFL KMNSLQIDDT AIYYCAREAS
101 RDHYFDYWGO GTTLTVSSAS TKGPSVFPLA PSSKSTSGGT AALGCLVKDY
151 FPEPVTVSWN SGALTSGVHT FPAVLQSSGL YSLSSVVTVP SSSLGTQTYI
201 CNVNHKPSNT KVDKRVEPKS CDKTHTCPPC PAPELLGGPS VFLFPPKPKD
251 TLMISRTPEV TCVVVDVSHE DPEVKFNWYV DGVEVHNAKT KPREEQYNST
301 YRVVSVLTVL HQDWLNGKEY KCKVSNKALP APIEKTISKA KGQPREPQVY
351 TLPPSREEMT KNQVSLTCLV KGFYPSDIAV EWESNGQPEN NYKTTTPVLD
401 SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH EALHNHYTQK SLSLSPGK

```

The variable regions are underlined.

The antibodies were expressed in 293EBNA monolayer cells. The plasmids were transfected using PEI based transfection reagents. The transfected cells were incubated in Opti-MEM serum free medium and the secreted antibodies were purified from medium using protein A/G affinity chromatography. The concentration of purified antibodies was determined by OD280 nm and the purity was measured by LabChip™ capillary electrophoresis.

The expression of both light and heavy chains was driven by human CMV promoter and bovine growth hormone polyadenylation signal. (Shiver *et al.*, *Ann. N.Y. Acad. Sci.*, 772:198-208, 1995.) The leader sequence in the front mediated the secretion of antibodies into the culture medium. The leader sequence for the heavy chain was MEWSWVFLFLSVTTGVHS (SEQ ID NO: 26) and for the light chain was MSVPTQVLGLLLWLTDARC (SEQ ID NO: 27). The expression vectors carry oriP from EBV viral genome for prolonged expression in 293EBNA cells and the bacterial sequences for kanamycin selection marker and replication origin in *E. coli*.

The antibodies were expressed in 293EBNA monolayer cells. The plasmids were transfected using PEI based transfection reagents. The transfected cells were incubated in Opti-MEM serum free medium and the secreted antibodies were purified from medium using protein A/G affinity chromatography. The concentration of purified antibodies was determined by OD280nm and the purity by LabChip capillary electrophoresis.

Example 11: Affinity Determination

Comparative analysis was performed on 2H2 mAb as hybridoma material, scAb and a chimeric antibody. 2H2 mAb V_H and V_L region were cloned and expressed as an IgG chimera as described in Example 10. scAb was cloned into the MP16 vector (Example 9), which produces a scFv with a Human Kappa chain tag fused to it. As further described below, the antigen affinity was not significantly different among the constructs.

To measure a 1:1 interaction between the binding domain and the antigen, the experimental set up on Biacore was modified depending on whether antibody fragment or full length IgG

was analyzed. For IgG measurements, the IgG was captured to the surface as ligand and ORF0657t was run as analyte. For antibody fragment analysis, ORF0657t was bound to the surface and the antibody fragment was run as the analyte. This demonstrated that the affinity of the original 2H2 mAb hybridoma material to the ORF0657t antigen shows no significant change upon recombinant cloning (Table 12). Data were acquired via surface plasmon resonance on a Biacore 3000; each analyte was run at multiple concentrations, with two replicates per concentration. Data were analyzed with BIAevaluation (Biacore, Inc.) with simultaneous fits of entire concentration series. Fit parameters are listed in Table 12.

Table 12

	On-rate k_a (1/Ms)	Off-rate k_d (1/s)	Affinity, KD	χ^2 global fit
2H2 murine IgG2b	6.10 E+04	2.01 E-03	33nM	0.902
2H2 scAb	4.91 E+04	1.91 E-03	39nM	0.429
2H2 IgG chimera	1.10 E+05	2.73 E-03	25nM	0.295

Example 12: ORF0657n Based Sequences

The highlighted amino acids (indicated by bold and underlying) present in SEQ ID NOs: 4-19 show amino acid alterations to ORF0657n:

0657n (SEQ ID NO: 1)

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKAVKPAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
 MKKENGEQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTPVKDYAYIRFSVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKSATIEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGDKDTPATKPTKGEVSSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

0657nC/e (SEQ ID NO: 2)

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKAVKPAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
 MKKENGEQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTPVKDYAYIRFSVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKSATIEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGDKDTPATKPTKGEVSSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKNLEHHHHHH

0657nH/y (SEQ ID NO: 3)

MAEETGGTNT~~EAQPKTE~~AVASPTTTSEKAPETKPVANAVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKA
TNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFEMKKKDGTTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQF
WRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGTKAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEE
DYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKAEYKKKLEDTKKALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDT
KYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMETTNDDYWKDFMVEGQVRV~~TISKDAKN~~NRTIIIFPYVEGKT
LYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKSNKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDD
NKQLPSVEKENDASSES~~SGDKT~~PATKPTKGEVSSSTPTTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSA
PLQKANIKNTNDGHTQSQNNKNTQENKAKS

SEQ ID NO: 4

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNT~~EAQPKTE~~AVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
MKKKDGTTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
EYKKKLEDTKKALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVME
TTNDDYWKDFMVEGQVRV~~TISKDAKN~~NRTIIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSES~~SGDKT~~PATKPTKGEVSSST
TPTTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 5

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNT~~EAQPKTE~~AVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPN~~WR~~PIDFE
MKKKDGTTQQFYHYASSVEPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
EYKKKLEDTKKALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVME
TTNDDYWKDFMVEGQVRV~~TISKDAKN~~NRTIIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSES~~SGDKT~~PATKPTKGEVSSST
TPTTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 6

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNT~~EAQPKTE~~AVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPN~~WR~~PIDFE
MKKKDGTTQQFYHYASSVEPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYEL~~EKI~~QDKLPEKLKA
EYKKKLEDTKKALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVME
TTNDDYWKDFMVEGQVRV~~TISKDAKN~~NRTIIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSES~~SGDKT~~PATKPTKGEVSSST
TPTTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 7

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTV KDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLEQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALAEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 8

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTV KDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLEQVYELEKI QDKLPEKLKA
 EYKKKLEDTKKALAEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 9

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTKSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTQDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLEQVYELEKI QDKLPEKLKA
 EYKKKLEDTKKALAEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 10

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTKSKPEIELGLQSGSTWRKFEVYEGDKKLPIKLVSYDTQDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLEQVYELEKI QDKLPEKLKA
 EYKKKLEDTKKALAEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST

TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLOKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 11

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
MKNDKGTQQFYHYASSVEPARVIFTKSKPIIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTDKDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTTLERQVYELEKIQDKLPEKLKA
EYKKKLEDTKKALAEQVKSATTEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKMYMME
TTNDDYWKDFMVEGQVRVITISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLOKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 12

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
MKNDKGTQQFYHYASSVEPARVIFTKSKPIIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTDKDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTTLERQVYELEKIQDKLPEKLKA
EYKKKLEQTKKALAEQVKSATTEFQNVQPTNEKMTDLQDAHYVVYESVENSESMDTFVKHPIKTGMLNGKMYMME
TTNDDYWKDFMVEGQVRVITISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLOKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 13

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
MKNDKGTQQFYHYASSVEPARVIFTKSKPIIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTDKDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTTLERQVYELEKIQDKLPEKLKA
EYKKKLEQTKKALAEQVKSATTEFQNVQPTNEKMTDLQDAHYVVYESVENSESMDTFVKHPIKTGMLNGKMYMME
TTNDDYWKDFMVEGKRVITISKDAKNNTRTIIFPYVEGKALYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLOKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPRKRKN

SEQ ID NO: 14

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKHSAPNSRPIDFE
MKKKDGTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVKDYAYIRFSVSNGT
KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTTLERQVYELNKLQEKLPEKLKA
EYKKKLEDTKKALDEQVKSAYTEFQNVQPTNDKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKMYMME

TTNDDYWKDFMVEGQSVRTISKDAKNNTRTIIFPYIEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDPTATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 15

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELRDAIKNPAIKDKHSAPNSRPIDFE
 MKKKDGTQQFYHYASTVKPARVIFTDTKPEIELGLQSGQFWRKFEVYEGDKKL~~P~~VKLVSYS~~D~~VKDYAYIRFSVNGT
~~R~~AVKIVSSTH~~N~~NNKEEKYDYTLMEFAQPIYNSADKYKTEEDYKAEKLLAPYKAKT~~L~~ERQVYELNK~~L~~QDKLPEKLKA
 EYKKKL~~D~~DTKKALDDQVKSAYTEFQNVQPTNEKMTDLQDTKYV~~V~~ESVENNESVMDTFVKHPIKTGMLNGKKY~~V~~VME
 TTNDDYWKDFIVEGQSVRTISKDAKNNTRTIIFPYIEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDPTATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 16

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAI~~I~~DKHSAPNSRPIDFE
 MKKKDGTQQFYHYASSVKPARVIFTDS~~G~~PEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVKDYAYIRF~~P~~VSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFTEEDYKAEKLLAPYKAKT~~L~~ERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKS~~A~~ITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKY~~M~~VME
 TTNDDYWKDFMVEGQSVRTISKDAKNNTRTIIFPYIEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDPTATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 17

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAI~~I~~DKHSAPNSRPIDFE
 MKKKDGTQQFYHYASSVKPARVIFTDS~~G~~PEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVKDYAYIRF~~P~~VSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFTEEDYKAEKLLAPYKAKT~~L~~ERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKS~~A~~ITEFQNVQPTNEKMTDLQDTKYVVYES~~E~~ENNESMMDTFVKHPI~~Y~~TGMLNGKKY~~M~~VME
 TTNDDYWKDFMVEGQSVRTISKDAKNNTRTIIFPYIEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDPTATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 18

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIIDKDHSAPNSRPIDFE
 MKKKDGTQQFYHYASSVKPARVIFTDSGPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFPVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKDEEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESEENNESMMDTFVKHPIYTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDPTPATKPTKGEVSSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 19

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREGSEAIKNPAIKDKHSAPNSRPI
 DFEMKKDGTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVS
 NGTKAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEK
 LKAEYKKKLEDTKKALDEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYM
 VMETTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVVDDKEAFTKA
 NTDKSNKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDPTPATKPTKGEV
 ESSSTTPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLP
 QTGEESNKDMTLPMLALLALSSIVAFVLPKRKN

Other embodiments are within the following claims. While several embodiments have been shown and described, various modifications may be made without departing from the spirit and scope of the present invention.

WHAT IS CLAIMED IS:

1. An isolated antigen binding protein comprising a first variable region and a second variable region, wherein said binding protein binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region.
2. The binding protein of claim 1, wherein said target region is the mAb 2H2.BE11 target region and said first variable region is a V_H region comprising:
 - a first V_H CDR comprising amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid;
 - a second V_H CDR comprising amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acid; and
 - a third V_H CDR comprising amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.
3. The binding protein of claim 2, wherein said second variable region is a V_L region comprising:
 - a first V_L CDR comprising amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid;
 - a second V_L CDR comprising amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and
 - a third V_L CDR comprising amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.
4. The binding protein of claim 3, wherein said binding protein is an antibody.
5. The binding protein of claim 4, wherein said antibody is a monoclonal antibody.
6. The binding protein of claim 4, wherein said V_H region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and said V_L region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.
7. The binding protein of claim 6, wherein said binding protein is an antibody comprising (a) a heavy chain comprising said V_H region, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄ subtype; and (b) a light chain comprising said V_L region, and either a human kappa C_L or human lambda C_L.

8. The binding protein of claim 3, wherein
said V_H region comprises said first V_H CDR consisting of amino acids 36-45 of SEQ ID NO: 20, said second V_H CDR consisting of amino acids 50-65 of SEQ ID NO: 20, and said third V_H CDR consisting of amino acids 98-107 of SEQ ID NO: 20 and;
said first V_L region comprises said first V_L CDR consisting of amino acids 24-33 of SEQ ID NO: 21, said second V_L CDR consisting of amino acids 49-55 of SEQ ID NO: 21, and said third V_L CDR consisting of amino acids 88-96 of SEQ ID NO: 21.

9. The binding protein of claim 8, wherein said binding protein is an antibody.

10. The binding protein of claim 9, wherein said antibody is a monoclonal antibody.

11. The binding protein of claim 9, wherein said V_H region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and said V_L region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

12. The binding protein of claim 8, wherein said binding protein is an antibody comprising (a) a heavy chain comprising said V_H region, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄ subtype; and (b) a light chain comprising said V_L region, and either a human kappa C_L or human lambda C_L.

13. The binding protein of claim 12, wherein said heavy chain consists essentially of the amino acid sequence of SEQ ID NO: 22; and said light chain consists essentially of the amino acid sequence of SEQ ID NO: 23.

14. A nucleic acid comprising a recombinant gene comprising a nucleotide sequence encoding an antibody variable region that binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region.

15. The nucleic acid of claim 14, wherein said target region is the mAb 2H2.BE11 target region and said variable region is a V_H region comprising:
a first V_H CDR comprising amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid;
a second V_H CDR comprising amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acids; and

a third V_h CDR comprising amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.

16. The nucleic acid of claim 15, wherein said V_h region comprises said first V_h CDR consisting of amino acids 36-45 of SEQ ID NO: 20; said second V_h CDR consisting of amino acids 50-65 of SEQ ID NO: 20; and said third V_h CDR consisting of amino acids 98-107 of SEQ ID NO: 20.

17. The nucleic acid of claim 14, wherein said variable region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20.

18. The nucleic acid of claim 17, wherein said recombinant gene encodes an antibody heavy chain comprising said variable region, a human hinge, and CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄ subtype.

19. The nucleic acid of claim 14, wherein said variable region is a V_l region comprising:
a first V_l CDR comprising amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid;
a second V_l CDR comprising amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and
a third V_l CDR comprising amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.

20. The nucleic acid of claim 19, wherein said first V_l region comprises said first V_l CDR consisting of amino acids 24-33 of SEQ ID NO: 21, said second V_l CDR consisting of amino acids 49-55 of SEQ ID NO: 21, and said third V_l CDR consisting of amino acids 88-96 of SEQ ID NO: 21.

21. The nucleic acid of claim 14, wherein said variable region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

22. The nucleic acid of claim 17, wherein said recombinant gene encodes an antibody light chain comprising said variable region and a human kappa or lambda C_L.

23. A recombinant cell comprising one or more nucleic acids of any one of claims 14-22.

24. The recombinant cell of claim 23, wherein said cell comprises both the nucleic acid of claim 18 and the nucleic acid of claim 22.

25. A method of producing protein comprising an antibody variable region comprising the steps of:

- a) growing the recombinant cell of claim 23 under conditions wherein said protein is expressed; and
- b) purifying said protein.

26. A method of producing protein comprising an antibody variable region comprising the steps of:

- a) growing the recombinant cell of claim 24 under conditions wherein said protein is expressed; and
- b) purifying said protein.

27. A pharmaceutical composition comprising the binding protein of any one of claims 1-13 and a pharmaceutically acceptable carrier.

28. A method of detecting the presence of an OFR0657n antigen in a solution or on a cell comprising the steps of: (a) providing the binding protein of any one of claims 1-13 to said solution or said cell; and (b) measuring the ability of said binding protein to bind to said antigen present in said solution or to said cell.

29. A method of treating against an *S. aureus* infection in a patient comprising the step of administering to said patient an effective amount of the binding protein of any one of claims 1-13.

30. The method of claim 29, wherein said antigen binding protein is administered in conjunction with surgery or a foreign body implant.

31. A cell line producing a protein that is either mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11, or that competes with either mAb 1G3.B3., mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11 for binding to ORF0657n.

32. The cell line of claim 31, wherein said line is either ATCC No: PTA-7124, ATCC No: PTA-7125, ATCC No: PTA-7126 or ATCC No: PTA-7127.

1/5





FIG. 1

2/5

**FIG. 2**

3/5




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4A

4/5




FIG. 4B




FIG. 5A





FIG. 5B

5/5

**FIG. 5C**

SEQUENCE LISTING

<110> Merck & Co., Inc.

<120> ANTIGEN-BINDING PROTEINS TARGETING S.
AUREUS ORF0657n

<130> 21934Y PCT

<150> 60/763,023

<151> 2006-01-27

<160> 34

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 645

<212> PRT

<213> Staphylococcus aureus

<400> 1

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1          5          10          15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu
 20          25          30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
 35          40          45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50          55          60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
 65          70          75          80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
 85          90          95
Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Ala Val Lys
100          105          110
Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
115          120          125
Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
130          135          140
Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Glu Asn Gly
145          150          155          160
Glu Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val
165          170          175
Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
180          185          190
Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
195          200          205
Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg
210          215          220
Phe Ser Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr
225          230          235          240
His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe
245          250          255

```


Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	
			260					265					270			
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	
		275					280					285				
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	
	290					295				300						
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala	
305					310					315					320	
Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	
			325						330					335		
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val	
			340					345					350			
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys	
		355					360					365				
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	
	370					375					380					
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	
385					390					395					400	
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile	
				405					410					415		
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys	
			420					425					430			
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile	
		435					440					445				
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys	
	450					455				460						
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr	
465					470					475					480	
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln	
				485					490					495		
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu	
			500					505					510			
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys	
		515					520					525				
Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val	
		530				535					540					
Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys	
545					550					555					560	
Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys	
			565						570					575		
Asp	Ser	Ala	Pro	Leu	Gln	Lys	Ala	Asn	Ile	Lys	Asn	Thr	Asn	Asp	Gly	
			580					585					590			
His	Thr	Gln	Ser	Gln	Asn	Asn	Lys	Asn	Thr	Gln	Glu	Asn	Lys	Ala	Lys	
		595					600					605				
Ser	Leu	Pro	Gln	Thr	Gly	Glu	Glu	Ser	Asn	Lys	Asp	Met	Thr	Leu	Pro	
	610					615					620					
Leu	Met	Ala	Leu	Leu	Ala	Leu	Ser	Ser	Ile	Val	Ala	Phe	Val	Leu	Pro	
625					630					635					640	
Arg	Lys	Arg	Lys	Asn												
				645												

<210> 2

<211> 653

<212> PRT

<213> Artificial Sequence

<220>

<223> 0657n with a carboxy His-Tag

<400> 2

Met	Asn	Lys	Gln	Gln	Lys	Glu	Phe	Lys	Ser	Phe	Tyr	Ser	Ile	Arg	Lys
1				5					10					15	
Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	
		20						25					30		
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr
		35					40					45			
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr
	50					55					60				
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser
65					70					75					80
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala
			85						90					95	
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Ala	Val	Lys
			100					105					110		
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu
		115					120					125			
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser
	130					135					140				
Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Glu	Asn	Gly
145					150					155					160
Glu	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Lys	Pro	Ala	Arg	Val
			165						170					175	
Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly
		180					185						190		
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
	195						200					205			
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
	210					215					220				
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Ala	Val	Lys	Ile	Val	Ser	Ser	Thr
225					230					235					240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe
			245						250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp
		260						265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
		275					280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala
305					310					315					320
Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
			325						330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
		340						345					350		
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
		355					360					365			
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln
385					390					395					400

```

Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
      405      410      415
Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
      420      425      430
Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
      435      440      445
Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
      450      455      460
Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
465      470      475      480
Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
      485      490      495
Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
      500      505      510
Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
      515      520      525
Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
      530      535      540
Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
545      550      555      560
Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
      565      570      575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
      580      585      590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
      595      600      605
Ser Leu Pro Gln Thr Gly Glu Ser Asn Lys Asp Met Thr Leu Pro
      610      615      620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
625      630      635      640
Arg Lys Arg Lys Asn Leu Glu His His His His His His
      645      650

```

<210> 3

<211> 569

<212> PRT

<213> Artificial Sequence

<220>

<223> Truncated version of ORF0657n

<400> 3

```

Met Ala Glu Glu Thr Gly Gly Thr Asn Thr Glu Ala Gln Pro Lys Thr
1      5      10      15
Glu Ala Val Ala Ser Pro Thr Thr Thr Ser Glu Lys Ala Pro Glu Thr
      20      25      30
Lys Pro Val Ala Asn Ala Val Ser Val Ser Asn Lys Glu Val Glu Ala
      35      40      45
Pro Thr Ser Glu Thr Lys Glu Ala Lys Glu Val Lys Glu Val Lys Ala
      50      55      60
Pro Lys Glu Thr Lys Glu Val Lys Pro Ala Ala Lys Ala Thr Asn Asn
65      70      75      80
Thr Tyr Pro Ile Leu Asn Gln Glu Leu Arg Glu Ala Ile Lys Asn Pro
      85      90      95

```

Ala	Ile	Lys	Asp	Lys	Asp	His	Ser	Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp
			100					105					110		
Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly	Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala
		115					120					125			
Ser	Ser	Val	Lys	Pro	Ala	Arg	Val	Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu
		130				135					140				
Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly	Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val
145					150					155				160	
Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro	Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr
			165						170					175	
Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg	Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys
		180						185					190		
Ala	Val	Lys	Ile	Val	Ser	Ser	Thr	His	Phe	Asn	Asn	Lys	Glu	Glu	Lys
		195					200					205			
Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe	Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala
		210				215					220				
Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala
225					230					235				240	
Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn
				245					250					255	
Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys
		260						265					270		
Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala	Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala
		275					280					285			
Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp
	290				295						300				
Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val	Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu
305					310					315					320
Ser	Met	Met	Asp	Thr	Phe	Val	Lys	His	Pro	Ile	Lys	Thr	Gly	Met	Leu
			325					330						335	
Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp
		340						345					350		
Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp
		355					360					365			
Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile	Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys
		370				375					380				
Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys	Val	His	Val	Lys	Thr	Ile	Asp	Tyr
385					390					395					400
Asp	Gly	Gln	Tyr	His	Val	Arg	Ile	Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys
			405					410						415	
Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys	Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala
		420						425					430		
Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr	Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro
		435					440					445			
Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln	Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys
		450				455					460				
Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu	Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly
465					470					475					480
Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys	Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser
			485						490					495	
Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val	Val	Ser	Thr	Thr	Gln	Asn	Val	Ala
		500						505					510		
Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys	Thr	Thr	Lys	Asp	Val	Val	Gln	Thr
		515					520					525			

Ser Ala Gly Ser Ser Glu Ala Lys Asp Ser Ala Pro Leu Gln Lys Ala
 530 535 540
 Asn Ile Lys Asn Thr Asn Asp Gly His Thr Gln Ser Gln Asn Asn Lys
 545 550 555 560
 Asn Thr Gln Glu Asn Lys Ala Lys Ser
 565

<210> 4
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

<400> 4
 Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1 5 10 15
 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu
 20 25 30
 Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
 35 40 45
 Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50 55 60
 Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
 65 70 75 80
 Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
 85 90 95
 Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
 100 105 110
 Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
 115 120 125
 Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
 130 135 140
 Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Lys Asp Gly
 145 150 155 160
 Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val
 165 170 175
 Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
 180 185 190
 Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
 195 200 205
 Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg
 210 215 220
 Phe Ser Val Ser Asn Gly Thr Lys Glu Val Lys Ile Val Ser Ser Thr
 225 230 235 240
 His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe
 245 250 255
 Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp
 260 265 270
 Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu
 275 280 285
 Glu Arg Gln Val Tyr Glu Leu Asn Lys Ile Gln Asp Lys Leu Pro Glu
 290 295 300

Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala
 305 310 315 320
 Leu Asp Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln
 325 330 335
 Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val
 340 345 350
 Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys
 355 360 365
 His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met
 370 375 380
 Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln
 385 390 395 400
 Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
 405 410 415
 Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
 420 425 430
 Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
 435 440 445
 Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
 450 455 460
 Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
 465 470 475 480
 Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
 485 490 495
 Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
 500 505 510
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515 520 525
 Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530 535 540
 Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545 550 555 560
 Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
 565 570 575
 Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
 580 585 590
 His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
 595 600 605
 Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
 610 615 620
 Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625 630 635 640
 Arg Lys Arg Lys Asn
 645

<210> 5

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 5

Met	Asn	Lys	Gln	Gln	Lys	Glu	Phe	Lys	Ser	Phe	Tyr	Ser	Ile	Arg	Lys	
1				5					10					15		
Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	Leu	
			20					25					30			
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr	
		35				40						45				
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr	
	50					55					60					
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser	
65					70					75					80	
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala	
				85					90					95		
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys	
			100					105					110			
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu	
		115					120					125				
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser	
	130					135					140					
Ala	Pro	Asn	Trp	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly	
145					150					155					160	
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Glu	Pro	Ala	Arg	Val	
				165					170					175		
Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly	
			180					185					190			
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro	
		195					200					205				
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg	
	210					215					220					
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Glu	Val	Lys	Ile	Val	Ser	Ser	Thr	
225					230					235					240	
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe	
				245					250					255		
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	
			260					265					270			
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	
		275					280					285				
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	
	290					295					300					
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala	
305					310					315					320	
Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	
				325					330					335		
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val	
		340						345					350			
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys	
		355					360					365				
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	
	370					375					380					
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	
385					390					395					400	
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile	
				405					410					415		
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys	
			420					425					430			

Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
 435 440 445
 Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
 450 455 460
 Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
 465 470 475 480
 Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
 485 490 495
 Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
 500 505 510
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515 520 525
 Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530 535 540
 Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545 550 555 560
 Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
 565 570 575
 Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
 580 585 590
 His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
 595 600 605
 Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
 610 615 620
 Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625 630 635 640
 Arg Lys Arg Lys Asn
 645

<210> 6
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

<400> 6
 Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1 5 10 15
 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
 20 25 30
 Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
 35 40 45
 Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50 55 60
 Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
 65 70 75 80
 Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
 85 90 95
 Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
 100 105 110
 Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
 115 120 125

Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
 130 135 140
 Ala Pro Asn Trp Arg Pro Ile Asp Phe Glu Met Lys Lys Lys Asp Gly
 145 150 155 160
 Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Glu Pro Ala Arg Val
 165 170 175
 Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
 180 185 190
 Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
 195 200 205
 Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg
 210 215 220
 Phe Ser Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr
 225 230 235 240
 His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe
 245 250 255
 Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp
 260 265 270
 Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu
 275 280 285
 Glu Arg Gln Val Tyr Glu Leu Glu Lys Ile Gln Asp Lys Leu Pro Glu
 290 295 300
 Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala
 305 310 315 320
 Leu Asp Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln
 325 330 335
 Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val
 340 345 350
 Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys
 355 360 365
 His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met
 370 375 380
 Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln
 385 390 395 400
 Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
 405 410 415
 Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
 420 425 430
 Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
 435 440 445
 Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
 450 455 460
 Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
 465 470 475 480
 Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
 485 490 495
 Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
 500 505 510
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515 520 525
 Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530 535 540
 Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545 550 555 560

```

Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
      565                               570                               575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
      580                               585                               590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
      595                               600                               605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
      610                               615                               620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
      625                               630                               635                               640
Arg Lys Arg Lys Asn
      645

```

<210> 7

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 7

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
1      5      10      15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu
      20      25      30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
      35      40      45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
50      55      60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
65      70      75      80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
      85      90      95
Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
100      105      110
Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
115      120      125
Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
130      135      140
Ala Pro Asn Trp Arg Pro Ile Asp Phe Glu Met Lys Lys Lys Asp Gly
145      150      155      160
Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Glu Pro Ala Arg Val
165      170      175
Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
180      185      190
Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
195      200      205
Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg
210      215      220
Phe Ser Val Ser Asn Gly Thr Lys Glu Val Lys Ile Val Ser Ser Thr
225      230      235      240
His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Val Phe
245      250      255

```

Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	
			260					265					270			
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	
		275					280					285				
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	
	290					295					300					
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala	
305					310					315					320	
Leu	Ala	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	
				325					330					335		
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val	
			340					345					350			
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys	
		355					360					365				
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	
	370					375						380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	
385					390					395					400	
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile	
				405					410					415		
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys	
			420					425					430			
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile	
		435					440					445				
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys	
	450					455					460					
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr	
465					470					475					480	
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln	
				485					490					495		
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu	
			500					505					510			
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys	
		515					520					525				
Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val	
						535						540				
Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys	
545					550					555					560	
Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys	
				565					570					575		
Asp	Ser	Ala	Pro	Leu	Gln	Lys	Ala	Asn	Ile	Lys	Asn	Thr	Asn	Asp	Gly	
			580					585					590			
His	Thr	Gln	Ser	Gln	Asn	Asn	Lys	Asn	Thr	Gln	Glu	Asn	Lys	Ala	Lys	
		595					600					605				
Ser	Leu	Pro	Gln	Thr	Gly	Glu	Glu	Ser	Asn	Lys	Asp	Met	Thr	Leu	Pro	
	610					615					620					
Leu	Met	Ala	Leu	Leu	Ala	Leu	Ser	Ser	Ile	Val	Ala	Phe	Val	Leu	Pro	
625					630					635					640	
Arg	Lys	Arg	Lys	Asn												
				645												

<210> 8

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 8

Met	Asn	Lys	Gln	Gln	Lys	Glu	Phe	Lys	Ser	Phe	Tyr	Ser	Ile	Arg	Lys	1	5	10	15
Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	Leu	20	25	30	
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr	35	40	45	
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr	50	55	60	
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser	65	70	75	80
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala	85	90	95	
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys	100	105	110	
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu	115	120	125	
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser	130	135	140	
Ala	Pro	Asn	Trp	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly	145	150	155	160
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Glu	Pro	Ala	Arg	Val	165	170	175	
Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly	180	185	190	
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro	195	200	205	
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg	210	215	220	
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Glu	Val	Lys	Ile	Val	Ser	Ser	Thr	225	230	235	240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe	245	250	255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	260	265	270	
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	275	280	285	
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Glu	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	290	295	300	
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala	305	310	315	320
Leu	Ala	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	325	330	335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val	340	345	350	
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys	355	360	365	
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	370	375	380	
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	385	390	395	400

```

Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
      405                      410                      415
Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
      420                      425                      430
Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
      435                      440                      445
Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
      450                      455                      460
Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
      465                      470                      475                      480
Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
      485                      490                      495
Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
      500                      505                      510
Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
      515                      520                      525
Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
      530                      535                      540
Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
      545                      550                      555                      560
Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
      565                      570                      575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
      580                      585                      590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
      595                      600                      605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
      610                      615                      620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
      625                      630                      635                      640
Arg Lys Arg Lys Asn
      645

```

<210> 9

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 9

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
  1                      5                      10                      15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
      20                      25                      30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
      35                      40                      45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
      50                      55                      60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
      65                      70                      75                      80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
      85                      90                      95

```

Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys
			100					105					110		
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu
		115					120					125			
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser
	130					135					140				
Ala	Pro	Asn	Trp	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly
145					150					155					160
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Glu	Pro	Ala	Arg	Val
			165					170						175	
Ile	Phe	Thr	Lys	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly
			180					185					190		
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
		195					200					205			
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Asp	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
	210					215					220				
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Glu	Val	Lys	Ile	Val	Ser	Ser	Thr
225					230					235					240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe
			245					250						255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
		275					280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Glu	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala
305					310					315					320
Leu	Ala	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
			325					330						335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
			340					345					350		
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
		355				360						365			
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375						380			
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln
385					390					395					400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
			405					410						415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420					425					430		
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
		435				440						445			
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys
	450					455					460				
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr
465					470					475					480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln
			485					490						495	
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu
			500					505					510		
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys
		515					520					525			

```

Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
  530                      535                      540
Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
  545                      550                      555                      560
Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
                      565                      570                      575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
                      580                      585                      590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
                      595                      600                      605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
  610                      615                      620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
  625                      630                      635                      640
Arg Lys Arg Lys Asn
                      645

```

<210> 10
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

```

<400> 10
Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
  1                      5                      10                      15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
  20                      25                      30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
  35                      40                      45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
  50                      55                      60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
  65                      70                      75                      80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
  85                      90                      95
Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
  100                      105                      110
Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
  115                      120                      125
Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
  130                      135                      140
Ala Pro Asn Trp Arg Pro Ile Asp Phe Glu Met Lys Lys Lys Asp Gly
  145                      150                      155                      160
Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Glu Pro Ala Arg Val
  165                      170                      175
Ile Phe Thr Lys Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
  180                      185                      190
Ser Thr Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
  195                      200                      205
Ile Lys Leu Val Ser Tyr Asp Thr Asp Lys Asp Tyr Ala Tyr Ile Arg
  210                      215                      220

```

Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Glu	Val	Lys	Ile	Val	Ser	Ser	Thr
225					230					235					240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Val	Phe
				245					250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
		275					280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Glu	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295				300					
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala
305					310					315					320
Leu	Ala	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
				325					330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
			340				345						350		
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
		355					360						365		
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln
385					390					395					400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
				405					410					415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420					425					430		
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
		435					440					445			
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys
	450					455					460				
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr
465					470					475					480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln
				485					490					495	
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu
			500					505					510		
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys
		515					520					525			
Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val
						535						540			
Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys
545					550					555					560
Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys
				565					570					575	
Asp	Ser	Ala	Pro	Leu	Gln	Lys	Ala	Asn	Ile	Lys	Asn	Thr	Asn	Asp	Gly
			580					585					590		
His	Thr	Gln	Ser	Gln	Asn	Asn	Lys	Asn	Thr	Gln	Glu	Asn	Lys	Ala	Lys
		595					600					605			
Ser	Leu	Pro	Gln	Thr	Gly	Glu	Glu	Ser	Asn	Lys	Asp	Met	Thr	Leu	Pro
	610					615					620				
Leu	Met	Ala	Leu	Leu	Ala	Leu	Ser	Ser	Ile	Val	Ala	Phe	Val	Leu	Pro
625					630					635					640
Arg	Lys	Arg	Lys	Asn											
				645											

<210> 11
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

<400> 11
 Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1 5 10 15
 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu
 20 25 30
 Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
 35 40 45
 Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50 55 60
 Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
 65 70 75 80
 Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
 85 90 95
 Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
 100 105 110
 Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
 115 120 125
 Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
 130 135 140
 Ala Pro Asn Trp Arg Pro Ile Asp Phe Glu Met Lys Asn Asp Lys Gly
 145 150 155 160
 Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Glu Pro Ala Arg Val
 165 170 175
 Ile Phe Thr Lys Ser Lys Pro Ile Ile Glu Leu Gly Leu Gln Ser Gly
 180 185 190
 Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
 195 200 205
 Ile Lys Leu Val Ser Tyr Asp Thr Asp Lys Asp Tyr Ala Tyr Ile Arg
 210 215 220
 Phe Ser Val Ser Asn Gly Thr Lys Glu Val Lys Ile Val Ser Ser Thr
 225 230 235 240
 His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Val Phe
 245 250 255
 Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp
 260 265 270
 Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu
 275 280 285
 Glu Arg Gln Val Tyr Glu Leu Glu Lys Ile Gln Asp Lys Leu Pro Glu
 290 295 300
 Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala
 305 310 315 320
 Leu Ala Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln
 325 330 335
 Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val
 340 345 350
 Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys
 355 360 365

```

His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met
   370                               375                 380
Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln
 385                               390                 395                 400
Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
   405                               410                 415
Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
   420                               425                 430
Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
   435                               440                 445
Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
 450                               455                 460
Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
 465                               470                 475                 480
Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
   485                               490                 495
Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
   500                               505                 510
Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515                               520                 525
Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530                               535                 540
Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545                               550                 555                 560
Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
   565                               570                 575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
 580                               585                 590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
 595                               600                 605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
 610                               615                 620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625                               630                 635                 640
Arg Lys Arg Lys Asn
   645

```

<210> 12.

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 12

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1                               5                 10                 15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
   20                               25                 30
Met Ser Asn Gly Glu Ala Gln Ala Ala Ala Glu Glu Thr Gly Gly Thr
   35                               40                 45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50                               55                 60

```

Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser	65	70	75	80
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala	85	90	95	
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys	100	105	110	
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu	115	120	125	
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser	130	135	140	
Ala	Pro	Asn	Trp	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Asn	Asp	Lys	Gly	145	150	155	160
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Glu	Pro	Ala	Arg	Val	165	170	175	
Ile	Phe	Thr	Lys	Ser	Lys	Pro	Ile	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly	180	185	190	
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro	195	200	205	
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Asp	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg	210	215	220	
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Glu	Val	Lys	Ile	Val	Ser	Ser	Thr	225	230	235	240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Val	Phe	245	250	255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	260	265	270	
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	275	280	285	
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Glu	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	290	295	300	
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Gln	Thr	Lys	Lys	Ala	305	310	315	320
Leu	Ala	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	325	330	335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Ala	His	Tyr	Val	Val	340	345	350	
Tyr	Glu	Ser	Val	Glu	Asn	Ser	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys	355	360	365	
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	370	375	380	
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	385	390	395	400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile	405	410	415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys	420	425	430	
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile	435	440	445	
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys	450	455	460	
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr	465	470	475	480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln	485	490	495	

Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
 500 505 510
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515 520 525
 Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530 535 540
 Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545 550 555 560
 Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
 565 570 575
 Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
 580 585 590
 His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
 595 600 605
 Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
 610 615 620
 Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625 630 635 640
 Arg Lys Arg Lys Asn
 645

<210> 13
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

<400> 13
 Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1 5 10 15
 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
 20 25 30
 Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
 35 40 45
 Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50 55 60
 Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
 65 70 75 80
 Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
 85 90 95
 Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
 100 105 110
 Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
 115 120 125
 Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Asp His Ser
 130 135 140
 Ala Pro Asn Trp Arg Pro Ile Asp Phe Glu Met Lys Asn Asp Lys Gly
 145 150 155 160
 Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Glu Pro Ala Arg Val
 165 170 175
 Ile Phe Thr Lys Ser Lys Pro Ile Ile Glu Leu Gly Leu Gln Ser Gly
 180 185 190

Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
	195						200					205			
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Asp	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
	210					215					220				
Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Glu	Val	Lys	Ile	Val	Ser	Ser	Thr
225					230					235					240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Val	Phe
				245					250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
	275						280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Glu	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Gln	Thr	Lys	Lys	Ala
305					310					315					320
Leu	Ala	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
				325					330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Ala	His	Tyr	Val	Val
			340					345					350		
Tyr	Glu	Ser	Val	Glu	Asn	Ser	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
	355						360					365			
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Lys
385					390					395					400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
				405					410					415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Ala	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420					425					430		
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
	435						440					445			
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys
	450					455					460				
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr
465					470					475					480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln
				485					490					495	
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu
			500					505					510		
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys
	515						520					525			
Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val
	530					535						540			
Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys
545					550					555					560
Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys
				565					570					575	
Asp	Ser	Ala	Pro	Leu	Gln	Lys	Ala	Asn	Ile	Lys	Asn	Thr	Asn	Asp	Gly
			580					585					590		
His	Thr	Gln	Ser	Gln	Asn	Asn	Lys	Asn	Thr	Gln	Glu	Asn	Lys	Ala	Lys
	595						600					605			
Ser	Leu	Pro	Gln	Thr	Gly	Glu	Glu	Ser	Asn	Lys	Asp	Met	Thr	Leu	Pro
	610					615					620				

Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625 630 635 640
 Arg Lys Arg Lys Asn
 645

<210> 14
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

<400> 14
 Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1 5 10 15
 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
 20 25 30
 Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
 35 40 45
 Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
 50 55 60
 Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
 65 70 75 80
 Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
 85 90 95
 Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
 100 105 110
 Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
 115 120 125
 Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Lys Asp Lys Glu His Ser
 130 135 140
 Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Lys Asp Gly
 145 150 155 160
 Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val
 165 170 175
 Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
 180 185 190
 Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
 195 200 205
 Val Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg
 210 215 220
 Phe Ser Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr
 225 230 235 240
 His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe
 245 250 255
 Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp
 260 265 270
 Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu
 275 280 285
 Glu Arg Gln Val Tyr Glu Leu Asn Lys Leu Gln Glu Lys Leu Pro Glu
 290 295 300
 Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala
 305 310 315 320

Leu Asp Glu Gln Val Lys Ser Ala Val Thr Glu Phe Gln Asn Val Gln
 325 330 335
 Pro Thr Asn Asp Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val
 340 345 350
 Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys
 355 360 365
 His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met
 370 375 380
 Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln
 385 390 395 400
 Ser Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
 405 410 415
 Ile Phe Pro Tyr Ile Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
 420 425 430
 Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
 435 440 445
 Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
 450 455 460
 Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
 465 470 475 480
 Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
 485 490 495
 Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
 500 505 510
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515 520 525
 Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530 535 540
 Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545 550 555 560
 Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
 565 570 575
 Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
 580 585 590
 His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
 595 600 605
 Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
 610 615 620
 Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625 630 635 640
 Arg Lys Arg Lys Asn
 645

<210> 15
 <211> 645
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ORF0657n mutant

<400> 15
 Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1 5 10 15

Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	Leu
			20					25					30		
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr
		35				40						45			
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr
		50				55					60				
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser
		65			70					75					80
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala
			85					90						95	
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys
			100					105					110		
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu
		115						120				125			
Leu	Arg	Asp	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Glu	His	Ser
		130				135					140				
Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly
		145			150					155					160
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Thr	Val	Lys	Pro	Ala	Arg	Val
			165						170					175	
Ile	Phe	Thr	Asp	Thr	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly
			180					185					190		
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
		195					200					205			
Val	Lys	Leu	Val	Ser	Tyr	Asp	Ser	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
		210				215					220				
Phe	Ser	Val	Ser	Asn	Gly	Thr	Arg	Ala	Val	Lys	Ile	Val	Ser	Ser	Thr
		225			230					235					240
His	Tyr	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe
			245						250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Tyr	Lys	Thr	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
		275					280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Leu	Gln	Asp	Lys	Leu	Pro	Glu
		290				295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Asp	Asp	Thr	Lys	Lys	Ala
		305			310					315					320
Leu	Asp	Asp	Gln	Val	Lys	Ser	Ala	Val	Thr	Glu	Phe	Gln	Asn	Val	Gln
			325						330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
			340					345					350		
Phe	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Val	Met	Asp	Thr	Phe	Val	Lys
		355					360					365			
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Val	Val	Met
		370				375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Ile	Val	Glu	Gly	Gln
		385			390					395					400
Arg	Val	Arg	Thr	Val	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
			405					410						415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420					425				430			
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
		435					440					445			


```

Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
  450                      455                      460
Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
  465                      470                      475                      480
Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
                      485                      490                      495
Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
  500                      505                      510
Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
  515                      520                      525
Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
  530                      535                      540
Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
  545                      550                      555                      560
Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
                      565                      570                      575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
  580                      585                      590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
  595                      600                      605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
  610                      615                      620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
  625                      630                      635                      640
Arg Lys Arg Lys Asn
                      645

```

<210> 16

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 16

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
  1                      5                      10                      15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
                      20                      25                      30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
  35                      40                      45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
  50                      55                      60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
  65                      70                      75                      80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
                      85                      90                      95
Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
                      100                      105                      110
Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
                      115                      120                      125
Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Ile Asp Lys Asp His Ser
  130                      135                      140

```

Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly
145					150					155					160
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Lys	Pro	Ala	Arg	Val
			165						170					175	
Ile	Phe	Thr	Asp	Ser	Gly	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly
			180					185					190		
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
		195					200					205			
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
	210					215					220				
Phe	Pro	Val	Ser	Asn	Gly	Thr	Lys	Ala	Val	Lys	Ile	Val	Ser	Ser	Thr
225					230					235					240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe
				245					250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
	275						280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala
305					310					315					320
Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
				325					330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
			340					345					350		
Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
	355						360					365			
His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln
385					390					395					400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
				405					410					415	
Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys
			420					425					430		
Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile
	435						440					445			
Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys
	450					455					460				
Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr
465					470					475					480
Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln
				485					490					495	
Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu
			500					505					510		
Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys
	515						520					525			
Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val
	530					535					540				
Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys
545					550					555					560
Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys
				565					570					575	

```

Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
      580      585      590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
      595      600      605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
      610      615      620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
      625      630      635      640
Arg Lys Arg Lys Asn
      645

```

<210> 17

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 17

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
  1      5      10      15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
      20      25      30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
      35      40      45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
      50      55      60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
      65      70      75      80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
      85      90      95
Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
      100      105      110
Pro Ala Ala Lys Ala Thr Asn Asn Thr Tyr Pro Ile Leu Asn Gln Glu
      115      120      125
Leu Arg Glu Ala Ile Lys Asn Pro Ala Ile Ile Asp Lys Asp His Ser
      130      135      140
Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Lys Asp Gly
      145      150      155      160
Thr Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val
      165      170      175
Ile Phe Thr Asp Ser Gly Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly
      180      185      190
Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro
      195      200      205
Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg
      210      215      220
Phe Pro Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr
      225      230      235      240
His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe
      245      250      255
Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp
      260      265      270

```

Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu
 275 280 285
 Glu Arg Gln Val Tyr Glu Leu Asn Lys Ile Gln Asp Lys Leu Pro Glu
 290 295 300
 Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala
 305 310 315 320
 Leu Asp Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln
 325 330 335
 Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val
 340 345 350
 Tyr Glu Ser Glu Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys
 355 360 365
 His Pro Ile Tyr Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met
 370 375 380
 Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln
 385 390 395 400
 Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile
 405 410 415
 Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
 420 425 430
 Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
 435 440 445
 Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
 450 455 460
 Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
 465 470 475 480
 Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
 485 490 495
 Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
 500 505 510
 Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
 515 520 525
 Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
 530 535 540
 Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
 545 550 555 560
 Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
 565 570 575
 Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
 580 585 590
 His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
 595 600 605
 Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
 610 615 620
 Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
 625 630 635 640
 Arg Lys Arg Lys Asn
 645

<210> 18

<211> 645

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 18

Met	Asn	Lys	Gln	Gln	Lys	Glu	Phe	Lys	Ser	Phe	Tyr	Ser	Ile	Arg	Lys
1				5					10					15	
Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	
			20					25					30		
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr
		35					40					45			
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr
	50					55					60				
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser
65					70					75					80
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala
			85						90					95	
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys
			100					105					110		
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu
		115						120				125			
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Ile	Asp	Lys	Asp	His	Ser
	130					135					140				
Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly
145					150					155					160
Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Lys	Pro	Ala	Arg	Val
				165						170				175	
Ile	Phe	Thr	Asp	Ser	Gly	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly
			180					185					190		
Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro
	195						200					205			
Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg
	210					215					220				
Phe	Pro	Val	Ser	Asn	Gly	Thr	Lys	Ala	Val	Lys	Ile	Val	Ser	Ser	Thr
225					230					235					240
His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe
				245					250					255	
Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Asp	Glu	Glu	Asp
			260					265					270		
Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu
		275					280					285			
Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu
	290					295					300				
Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala
305					310					315					320
Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln
				325					330					335	
Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val
		340						345					350		
Tyr	Glu	Ser	Glu	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys
		355					360					365			
His	Pro	Ile	Tyr	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met
	370					375					380				
Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln
385					390					395					400
Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile
				405					410					415	

```

Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys
      420                      425      430
Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile
      435                      440      445
Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys
      450                      455      460
Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr
465      470                      475      480
Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln
      485                      490      495
Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu
      500                      505      510
Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys
      515                      520      525
Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val
      530                      535      540
Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys
545      550                      555      560
Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys
      565                      570      575
Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys Asn Thr Asn Asp Gly
      580                      585      590
His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln Glu Asn Lys Ala Lys
      595                      600      605
Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys Asp Met Thr Leu Pro
610      615                      620
Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val Ala Phe Val Leu Pro
625      630                      635      640
Arg Lys Arg Lys Asn
      645

```

<210> 19

<211> 650

<212> PRT

<213> Artificial Sequence

<220>

<223> ORF0657n mutant

<400> 19

```

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
 1      5      10      15
Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
      20      25      30
Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
      35      40      45
Asn Thr Glu Ala Gln Pro Lys Thr Glu Ala Val Ala Ser Pro Thr Thr
      50      55      60
Thr Ser Glu Lys Ala Pro Glu Thr Lys Pro Val Ala Asn Ala Val Ser
65      70      75      80
Val Ser Asn Lys Glu Val Glu Ala Pro Thr Ser Glu Thr Lys Glu Ala
      85      90      95
Lys Glu Val Lys Glu Val Lys Ala Pro Lys Glu Thr Lys Glu Val Lys
      100      105      110

```

Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu		
	115						120					125					
Leu	Arg	Glu	Gly	Ser	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys		
	130						135				140						
Asp	His	Ser	Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys		
	145				150					155					160		
Lys	Asp	Gly	Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Lys	Pro		
				165					170					175			
Ala	Arg	Val	Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu		
			180					185					190				
Gln	Ser	Gly	Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys		
	195						200					205					
Lys	Leu	Pro	Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala		
	210					215					220						
Tyr	Ile	Arg	Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Ala	Val	Lys	Ile	Val		
	225				230					235					240		
Ser	Ser	Thr	His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu		
				245					250					255			
Met	Glu	Phe	Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr		
			260					265					270				
Glu	Glu	Asp	Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala		
		275					280					285					
Lys	Thr	Leu	Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys		
	290					295					300						
Leu	Pro	Glu	Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr		
	305				310					315					320		
Lys	Lys	Ala	Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln		
				325					330					335			
Asn	Val	Gln	Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys		
			340					345					350				
Tyr	Val	Val	Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr		
		355					360					365					
Phe	Val	Lys	His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr		
	370					375					380						
Met	Val	Met	Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val		
	385				390					395					400		
Glu	Gly	Gln	Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr		
				405					410					415			
Arg	Thr	Ile	Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala		
			420					425					430				
Ile	Val	Lys	Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His		
		435					440					445					
Val	Arg	Ile	Val	Asp	Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr		
	450					455					460						
Asp	Lys	Ser	Asn	Lys	Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu		
	465				470					475					480		
Ala	Thr	Pro	Ala	Thr	Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys		
				485					490					495			
Glu	Ser	Gln	Lys	Gln	Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro		
		500						505					510				
Ser	Val	Glu	Lys	Glu	Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys		
		515					520					525					
Thr	Pro	Ala	Thr	Lys	Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr		
	530					535					540						

```

Thr Pro Thr Lys Val Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr
545          550          555          560
Thr Ala Ser Ser Lys Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly
          565          570          575
Ser Ser Glu Ala Lys Asp Ser Ala Pro Leu Gln Lys Ala Asn Ile Lys
          580          585          590
Asn Thr Asn Asp Gly His Thr Gln Ser Gln Asn Asn Lys Asn Thr Gln
          595          600          605
Glu Asn Lys Ala Lys Ser Leu Pro Gln Thr Gly Glu Glu Ser Asn Lys
          610          615          620
Asp Met Thr Leu Pro Leu Met Ala Leu Leu Ala Leu Ser Ser Ile Val
625          630          635          640
Ala Phe Val Leu Pro Arg Lys Arg Lys Asn
          645          650

```

<210> 20
 <211> 118
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 2H2 Vh

```

<400> 20
Asp Val His Leu Val Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
 1          5          10          15
Asn Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Arg Tyr
          20          25          30
Gly Val His Trp Val Arg Gln Pro Gly Lys Gly Leu Glu Trp Leu
          35          40          45
Gly Leu Ile Trp Ala Gly Gly Val Thr Ile Tyr Asn Ser Thr Leu Met
          50          55          60
Ser Arg Leu Ser Ile Ser Lys Asp Ser Ser Lys Ser Gln Val Phe Leu
65          70          75          80
Lys Met Asn Ser Leu Gln Ile Asp Asp Thr Ala Ile Tyr Tyr Cys Ala
          85          90          95
Arg Glu Ala Ser Arg Asp His Tyr Phe Asp Tyr Trp Gly Gln Gly Thr
          100          105          110
Thr Leu Thr Val Ser Ser
          115

```

<210> 21
 <211> 107
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 2H2 V1

```

<400> 21
Asp Ile Val Met Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1          5          10          15
Glu Lys Ile Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Ile
          20          25          30

```



```

Tyr Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
      35              40              45
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Phe Arg Phe Ser Gly Gly
      50              55              60
Gly Ser Gly Thr Ser Phe Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
65      70              75              80
Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
      85              90              95
Phe Gly Ala Gly Thr Lys Leu Glu Ile Lys Arg
      100              105

```

<210> 22
 <211> 213
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Mouse 2H2B3 Variable and Human Kappa Constant
 Region

```

<400> 22
Asp Ile Val Met Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1      5              10              15
Glu Lys Ile Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Ile
      20              25              30
Tyr Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
      35              40              45
Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Phe Arg Phe Ser Gly Gly
50      55              60
Gly Ser Gly Thr Ser Phe Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
65      70              75              80
Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
      85              90              95
Phe Gly Ala Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
      100              105              110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
      115              120              125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130      135              140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145      150              155              160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
      165              170              175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
      180              185              190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
195      200              205
Asn Arg Gly Glu Cys
      210

```

<210> 23
 <211> 448
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Mouse 2H2B3 Variable and Human IgG1 Constant
Region

<400> 23

Asp	Val	His	Leu	Val	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1				5					10					15	
Asn	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Arg	Tyr
			20					25					30		
Gly	Val	His	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu
		35				40						45			
Gly	Leu	Ile	Trp	Ala	Gly	Gly	Val	Thr	Ile	Tyr	Asn	Ser	Thr	Leu	Met
	50					55					60				
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu
65				70						75				80	
Lys	Met	Asn	Ser	Leu	Gln	Ile	Asp	Asp	Thr	Ala	Ile	Tyr	Tyr	Cys	Ala
			85						90					95	
Arg	Glu	Ala	Ser	Arg	Asp	His	Tyr	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
			100					105					110		
Thr	Leu	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro
		115					120					125			
Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly
	130					135					140				
Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn
145					150					155				160	
Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln
			165						170					175	
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser
			180					185					190		
Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser
	195						200					205			
Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr
	210					215					220				
His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser
225					230					235				240	
Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg
			245						250					255	
Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro
			260					265					270		
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala
		275					280					285			
Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val
	290					295					300				
Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr
305					310					315				320	
Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr
			325						330					335	
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu
			340					345					350		
Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys
		355					360					365			
Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser
	370					375					380				
Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp
385					390					395					400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 405 410 415
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 420 425 430
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435 440 445

<210> 24
 <211> 354
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Sequence encoding 2H2 Vh

<400> 24
 gatgtgcacc tgggtggagtc aggacctggc ctggtggcgc cctcacagaa cctgtccatc 60
 acttgcaactg tctctgggtt ctcattatcc cgctatgggtg tacactgggt tgcgcagcct 120
 ccaggaaagg gtctggagtg gctgggacta atatgggctg gtggagtcac aatttataat 180
 tcgactctca tgtccagact gagcatcagc aaagacagct ccaagagcca ggttttccta 240
 aaaatgaaca gtctacaaat tgatgacaca gccatttact actgtgccag agaagcatct 300
 cgggaccact actttgacta ctggggccaa ggcaccactc tcacagtctc ctcg 354

<210> 25
 <211> 321
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Sequence encoding 2H2 V1

<400> 25
 gatattgtga tgaccagtc tccagcaatc atgtctgcat ctccagggga gaagatcacc 60
 atgacctgca gtgccagctc aagtgttagt tacatctact ggtaccagca gaagtcaggc 120
 acctccccc aaagatggat ttatgacaca tccaaactgg cttctggagt cccttttcgc 180
 ttacagtggcg gtgggtctgg gacctctttc tctctcacia tcagcagcat ggaggtgaa 240
 gatgctgcca cttattactg ccagcagtggt agtagtaacc cactcacgtt cgggtgctggg 300
 accaagctgg aaataaaacg t 321

<210> 26
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Heavy chain leader sequence

<400> 26
 Met Glu Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly
 1 5 10 15
 Val His Ser

<210> 27
 <211> 20

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Light chain leader sequence

<400> 27
 Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Leu Trp Leu Thr
 1 5 10 15
 Asp Ala Arg Cys
 20

<210> 28
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<400> 28
 acagatgccca gatgcatat tgtgatgacc cagtct 36

<210> 29
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<400> 29
 tgcagccacc gtacgtttta tttccagctt ggtccc 36

<210> 30
 <211> 36
 <212> DNA
 <213> Oligonucleotide Primer

<400> 30
 acaggtgtcc actcggtatgt gcacctggtg gactca 36

<210> 31
 <211> 36
 <212> DNA
 <213> Oligonucleotide Primer

<400> 31
 gcccttggtg gatgccgagg agactgtgag agtggt 36

<210> 32
 <211> 20
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Linker

<400> 32

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10 15
Gly Gly Gly Ser
20

<210> 33

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<400> 33

gtattaggaa ttcggccccc gaggccgagg 30

<210> 34

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<400> 34

gcattactcg cggcccagcc ggccatggcg gac 33

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2007 (09.08.2007)

PCT

(10) International Publication Number
WO 2007/089470 A3

(51) International Patent Classification:

G01N 33/53 (2006.01) C12N 5/16 (2006.01)
C12N 5/06 (2006.01)

(21) International Application Number:

PCT/US2007/001687

(22) International Filing Date: 23 January 2007 (23.01.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/763,023 27 January 2006 (27.01.2006) US

(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BROWN, Martha, J. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). ANDERSON, Annaliesa, S. [GB/US];


126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). COPE, Leslie, D. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). JANSEN, Kathrin, Ute [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). MCNEELY, Tessie [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). HARVEY, Barrett [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). DURR, Erberhard [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). ERNST, Robin [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY,

[Continued on next page]

(54) Title: ANTIGEN-BINDING PROTEINS TARGETING S. AUREUS ORF0657N



(57) Abstract: The present invention features antigen binding protein that bind an ORF0657n target region (SEQ ID NO: 1). ORF0657n is an *S. aureus* protein. ORF0657n target regions are provided by the mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 binding sites. In a lethal model challenge, mAb 2H2.BE11 and mAb 13C7.BC1 provided for increased survival against *S. aureus* infection. There was also protection demonstrated in an *ex vivo* model with either the IgG1 or the IgG2b form of mAb 2H2; and in a passive immunization murine indwelling catheter model using mAb 2H2.BE11.



WO 2007/089470 A3



MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report*

- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

- *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

- (88) **Date of publication of the international search report:**
24 January 2008

(15) **Information about Correction:**

Previous Correction:

see PCT Gazette No. 44/2007 of 1 November 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 07/01687

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - G01N 33/53; C12N 5/06, 5/16 (2007.01)

USPC - 435/7.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC: 435/7.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC: 435/7.2, 70.21, 188.5, 326-332, 345 (see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST (DB=PGPB,USPT,EPAB,JPAB), PubMed, Google, WIPO

Terms: ORF0657, antibody, Staphylococcus aureus, staphylococcal, monoclonal
mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, mAb 13G11.BF3, hydridoma cell

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	WO 2005/009379 A2 (Anderson et al.) 03 February 2005 (03.02.2005), especially description, para [117]-[118]	1, 27-32 2-13
X — Y	US 6,979,446 B2 (Patti et al.) 27 December 2005 (27.12.2005), col 25, ln 44 to col 26, ln 1-6; col 4, ln 28-38; col 17, ln 65 to col 18, ln 34-44	14-18, 23, 25 2-13, 19-22
Y	US 6,806,079 B1 (McCafferty et al.) October 19, 2004 (10.19.2004), pg 223,-224, SEQ ID NO 243	3-13, 19-22

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 August 2007 (06.08.2007)

Date of mailing of the international search report

21 NOV 2007

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 07/01687

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 24 and 26
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2007 (09.08.2007)

PCT

(10) International Publication Number
WO 2007/089470 A2

(51) International Patent Classification:
A61K 39/40 (2006.01)

(21) International Application Number:
PCT/US2007/001687

(22) International Filing Date: 23 January 2007 (23.01.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/763,023 27 January 2006 (27.01.2006) US

(71) Applicant (for all designated States except US): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BROWN, Martha, J.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **ANDERSON, Annaliesa, S.** [GB/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **COPE, Leslie, D.** [US/US]; 126 East Lincoln


Avenue, Rahway, New Jersey 07065-0907 (US). **JANSEN, Kathrin, Ute** [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **MCNEELY, Tessie** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **HARVEY, Barrett** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **DURR, Erberhard** [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **ERNST, Robin** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS,

[Continued on next page]

(54) Title: ANTIGEN-BINDING PROTEINS TARGETING S. AUREUS ORF0657N



(57) Abstract: The present invention features antigen binding protein that bind an ORF0657n target region (SEQ ID NO: 1). ORF0657n is an *S. aureus* protein. ORF0657n target regions are provided by the mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 binding sites. In a lethal model challenge, mAb 2H2.BE11 and mAb 13C7.BC1 provided for increased survival against *S. aureus* infection. There was also protection demonstrated in an *ex vivo* model with either the IgG1 or the IgG2b form of mAb 2H2; and in a passive immunization murine indwelling catheter model using mAb 2H2.BE11.

WO 2007/089470 A2



RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *without international search report and to be republished upon receipt of that report*

— *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

- (48) **Date of publication of this corrected version:**

1 November 2007

- (15) **Information about Correction:**

see PCT Gazette No. 44/2007 of 1 November 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE OF THE INVENTION

ANTIGEN-BINDING PROTEINS TARGETING *S. AUREUS* ORF0657n

RELATED APPLICATIONS

- 5 The present application claims priority to U.S. Provisional Application No. 60/763,023, filed January 27, 2006, which is hereby incorporated by reference herein.

BACKGROUND OF THE INVENTION

- 10 The references cited throughout the present application are not admitted to be prior art to the claimed invention.

- Staphylococcus aureus* is a pathogen responsible for a wide range of diseases and conditions. Examples of diseases and conditions caused by *S. aureus* include bacteremia, infective endocarditis, folliculitis, furuncle, carbuncle, impetigo, bullous impetigo, cellulitis, botryomycosis, toxic shock syndrome, scalded skin syndrome, central nervous system infections, infective and inflammatory eye disease, osteomyelitis and other infections of joints and bones, and respiratory tract infections. (*The Staphylococci in Human Disease*, Crossley and Archer (eds.), Churchill Livingstone Inc. 1997.)

- 15 Immunological based strategies can be employed to control *S. aureus* infections and the spread of *S. aureus*. Immunological based strategies include passive and active immunization. Passive immunization employs immunoglobulins targeting *S. aureus*. Active immunization induces immune
20 responses against *S. aureus*.

SUMMARY OF THE INVENTION

- The present invention features antigen binding protein that bind an ORF0657n target region (SEQ ID NO: 1). ORF0657n is an *S. aureus* protein. ORF0657n target regions are provided by the
25 mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 binding sites. In a lethal model challenge, mAb 2H2.BE11 and mAb 13C7.BC1 provided for increased survival against *S. aureus* infection. There was also protection demonstrated in an *ex vivo* model with either the IgG1 or the IgG2b form of mAb 2H2; and in a passive immunization murine indwelling catheter model using mAb 2H2.BE11.

- 30 Mouse hybridoma cell lines producing mAb 1G3.BD4, mAb 2H2.BE11; mAb 13C7.BC1, and mAb 13G11.BF3 were deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, in accordance with Budapest Treaty on September 30, 2005. The cells lines were designated: ATCC No. PTA-7124 (producing mAb 2H2.BE11), ATCC No. PTA-7125 (producing mAb 13C7.BC1), ATCC No. PTA-7126 (producing mAb 1G3.BD4), and ATCC
35 No. PTA-7127 (producing mAb 13G11.BF3).

 Thus, a first aspect of the present invention features an isolated antigen binding protein comprising a first variable region and a second variable region. The first and second variable regions

bind one or more target regions selected from the group consisting of: mAb 1G3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1 target region, and mAb 13G11.BF3 target region.

Reference to "isolated" indicates a different form than found in nature. The different form can be, for example, a different purity than found in nature and/or a structure that is not found in nature. A structure not found in nature includes recombinant structures where different regions are combined together, for example, humanized antibodies where one or more murine complementary determining regions is inserted onto a human framework scaffold or a murine antibody is resurfaced to resemble the surface residues of a human antibody, hybrid antibodies where one or more complementary determining regions from an antigen binding protein is inserted into a different framework scaffold, and antibodies derived from natural human sequences where genes coding for light and heavy variable domains were randomly combined together.

The isolated protein is preferably substantially free of serum proteins. A protein substantially free of serum proteins is present in an environment lacking most or all serum proteins.

A "variable region" has the structure of an antibody variable region from a heavy or light chain. Antibody heavy and light chain variable regions contain three complementary determining regions interspaced onto a framework. The complementary determining regions are primarily responsible for recognizing a particular epitope.

A target region is defined with respect to the ORF0657n region (SEQ ID NO: 1) bound by mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, or mAb 13G11.BF3. For example, the mAb 1G3.BD4 target region is the ORF0657n region to which mAb 1G3.BD4 binds.

A protein binding an identified target region competes with either mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, or mAb 13G11.BF3 for binding to the target region. For example, a protein competing with mAb 1G3.BD4 binding to ORF0657n binds to the mAb 1G3.BD4 target region.

A protein that competes with either the monoclonal antibody mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11 reduces binding of the monoclonal antibody to ORF0657n by at least about 20%, preferably at least about 50%, when excess and equal amounts of the competing protein and monoclonal antibody are employed.

Reference to "protein" indicates a contiguous amino acid sequence and does not provide a minimum or maximum size limitation. One or more amino acids present in the protein may contain a post-translational modification, such as glycosylation or disulfide bond formation.

A preferred antigen binding protein is a monoclonal antibody. Reference to a "monoclonal antibody" indicates a collection of antibodies having the same, or substantially the same, complementary determining region, and binding specificity. The variation in the antibodies is that which would occur if the antibodies were produced from the same construct(s).

Monoclonal antibodies can be produced, for example, from a particular hybridoma and from a recombinant cell containing one or more recombinant genes encoding the antibody. The antibody

may be encoded by more than one recombinant gene where, for example, one gene encodes the heavy chain and one gene encodes the light chain.

Another aspect of the present invention describes a nucleic acid containing a recombinant gene comprising a nucleotide sequence encoding an antibody variable region. The antibody
5 variable region can bind a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region.

A recombinant gene contains recombinant nucleic acid encoding a protein along with regulatory elements for proper transcription and processing (which may include translational and post translational elements). The recombinant nucleic acid by virtue of its sequence and/or form does not
10 occur in nature. Examples of recombinant nucleic acid include purified nucleic acid, two or more nucleic acid regions combined together providing a different nucleic acid than found in nature, and the absence of one or more nucleic acid regions (e.g., upstream or downstream regions) that are naturally associated with each other.

Another aspect of the present invention describes a recombinant cell comprising one or
15 more recombinant genes encoding an antibody variable region that binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region. Multiple recombinant genes are useful, for example, where one gene encodes an antibody heavy chain or fragment thereof containing the V_H region and another nucleic acid encodes an antibody light chain or fragment thereof containing the V_L region.

Another aspect of the present invention comprises a method of producing a protein
20 comprising an antibody variable region. The method comprising the steps of: (a) growing a recombinant cell comprising recombinant nucleotide acid encoding for a protein under conditions wherein the protein is expressed; and (b) purifying the protein.

Another aspect of the present invention describes a pharmaceutical composition. The
25 composition contains a therapeutically effective amount of an antigen binding protein and a pharmaceutically acceptable carrier.

A therapeutically effective amount is an amount sufficient to provide a useful therapeutic or prophylactic effect. For a patient infected with *S. aureus*, an effective amount is sufficient to achieve one or more of the following effects: reduce the ability of *S. aureus* to propagate in the patient or reduce
30 the amount of *S. aureus* in the patient. For a patient not infected with *S. aureus*, an effective amount is sufficient to achieve one or more of the following: a reduced susceptibility to *S. aureus* infection or a reduced ability of the infecting bacterium to establish persistent infection for chronic disease.

Another aspect of the present invention describes a method of detecting the presence of an OFR0657n antigen in a solution or on a cell. The method involves providing a binding protein
35 described herein to the solution or cell and measuring the ability of the binding protein to bind to the antigen in the solution or cell. Measurements can be quantitative or qualitative.

Reference to ORF0657n antigen includes full-length ORF0657n or a derivative thereof having an epitope that is recognized by mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11. Examples of derivatives include truncated versions; and full-length or truncated versions of ORF0657n containing one or more of the following amino acid alterations: one or more additions, one or more substitutions, and one or more deletions.

Another aspect of the present invention features a method of treating a patient against a *S. aureus* infection. The method comprises the step of administering to the patient an effective amount of an antigen binding protein described herein. The patient being treated may, or may not, be infected with *S. aureus*. Preferably, the patient is a human.

Another aspect of the present invention describes a cell line producing a protein that is either mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11, or that competes with either mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11 for binding to ORF0657n. Preferred cells lines are hybridomas, and recombinant cell lines containing recombinant nucleic acid encoding the protein.

Reference to open-ended terms such as "comprises" allows for additional elements or steps. Occasionally phrases such as "one or more" are used with or without open-ended terms to highlight the possibility of additional elements or steps.

Unless explicitly stated reference to terms such as "a" or "an" is not limited to one. For example, "a cell" does not exclude "cells". Occasionally phrases such as one or more are used to highlight the possible presence of a plurality.

Other features and advantages of the present invention are apparent from the additional descriptions provided herein including the different examples. The provided examples illustrate different components and methodology useful in practicing the present invention. The examples do not limit the claimed invention. Based on the present disclosure the skilled artisan can identify and employ other components and methodology useful for practicing the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the structure of an IgG molecule. "V_L" refers to a light chain variable region. "V_H" refers to a heavy chain variable region. "C_L" refers to a light chain constant region. "CH₁", "CH₂" and "CH₃" are heavy chain constant regions. Dashed lines indicate disulfide bonds.

Figure 2 illustrates a matrix outlining the reactivities of different monoclonal antibodies in a pair-wise binding study. The panel of monoclonal antibodies fell into three reactive areas by the BIACORE® method.

Figures 3A-3C: Groups of BALB/c mice (n = 20) were treated 20 hours prior to bacterial challenge with an i.p. injection of: ■, mAb 13C7.BC1; □, mAb 6G6.A8 (isotype control); or ○, PBS. Mice were challenged with *S. aureus* by i.v. injection and survival was monitored. Fig. 3A-

0.49 mg mAb 13C7.BC1; 0.45 mg mAb 6G6.A8; and 9.8×10^8 CFU *S. aureus* Becker. Fig. 3B- 0.49 mg mAb 13C7.BC1; 0.45 mg mAb 6G6.A8; and 9.6×10^8 CFU *S. aureus* Becker. Fig. 3C- 0.50 mg mAb 13C7.BC1; 0.45 mg mAb 6G6; and 9.9×10^8 CFU *S. aureus* Becker.

Figures 4A and 4B: Groups of BALB/c mice ($n = 20$) were treated 20 hours prior to bacterial challenge with an i.p. injection of: ■, mAb 13C7.BC1 (0.5 mg); □, mAb 6G6.A8 (isotype control) (0.5 mg); or ○, PBS (0.5 ml). Mice were challenged with *S. aureus* by i.v. injection and survival was monitored. Fig. 4A illustrates results with 2.09×10^8 CFU *S. aureus* UK58. Fig. 4B illustrates results with 2.15×10^8 *S. aureus* UK 58.

Figures 5A-5C: Groups of BALB/c mice ($n = 20$) were treated 20 hours prior to bacterial challenge with an i.p. injection of: ■, mAb 2H2.BE11, □, mAb 6G6.A8 (isotype control); ○, PBS. Mice were challenged with *S. aureus* by i.v. injection and survival was monitored. Fig. 5A- 0.43 mg mAb 2H2.BE11; 0.5 mg mAb 6G6.A8; and 9.8×10^8 CFU *S. aureus* Becker. Fig. 5B- 0.43 mg mAb 2H2.BE11; 0.5 mg mAb 6G6.A8; and 8.3×10^8 CFU *S. aureus* Becker. Fig. 5C- 0.43 mg mAb 2H2.BE11; 0.5 mg mAb 6G6.A8; and 9.3×10^8 CFU *S. aureus* Becker.

DETAILED DESCRIPTION OF THE INVENTION

ORF0657n is an *S. aureus* protein located at the *S. aureus* outer membrane. ORF0657n has been found to be well conserved in different strains of *S. aureus*. (Anderson *et al.*, International Publication No. WO 2005/009379, International Publication Date February 3, 2005.) Different ORF0657n derivatives can be used to produce a protective immune response against *S. aureus* infection. (Anderson *et al.*, International Publication No. WO 2005/009379, International Publication Date February 3, 2005.)

Due to their ability to recognize ORF0657n, the antigen binding proteins described herein can be used, for example, as a tool in the production, characterization, or study of ORF0657n based antigens. Antigen binding protein recognizing appropriate ORF0657n epitopes can also be used agent to treat *S. aureus* infection.

I. Antigen Binding Protein

Antigen binding proteins contain an antibody variable region providing for specific binding to an epitope. The antibody variable region can be present in, for example, a complete antibody, an antibody fragment, and a recombinant derivative of an antibody or antibody fragment.

Different classes of antibodies have different structures. Different antibody regions can be illustrated by reference to IgG (Figure 1). An IgG molecule contains four amino acid chains: two longer length heavy chains and two shorter light chains. The heavy and light chains each contain a constant region and a variable region. Within the variable regions are three hypervariable regions responsible for antigen specificity. (See, for example, Breitling *et al.*, Recombinant Antibodies, John Wiley & Sons, Inc. and Spektrum Akademischer Verlag, 1999; and Lewin, Genes IV, Oxford University Press and Cell Press, 1990.)

The hypervariable regions (also referred to as complementarity determining regions), are interposed between more conserved flanking regions (also referred to as framework regions). Amino acids associated with framework regions and complementarity determining regions can be numbered and aligned as described by Kabat *et al.*, Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services, 1991.

The two heavy chain carboxyl regions are constant regions joined by disulfide binding to produce an Fc region. The Fc region is important for providing antibody biological activity such as complement and macrophage activation. Each of the two heavy chains making up the Fc region extend into different Fab regions through a hinge region.

In higher vertebrates there are two classes of light chains and five classes of heavy chains. The light chains are either κ or λ . The heavy chains define the antibody class and are either α , δ , ϵ , γ , or μ . For example, IgG has a γ heavy chain. Subclasses also exist for different types of heavy chains such as human γ_1 , γ_2 , γ_3 , and γ_4 . Heavy chains impart a distinctive conformation to hinge and tail regions. (Lewin, Genes IV, Oxford University Press and Cell Press, 1990.)

Antibody fragments containing an antibody variable region include Fv, Fab, and Fab₂ regions. Each Fab region contains a light chain made up of a variable region and a constant region, and a heavy chain region containing a variable region and a constant region. A light chain is joined to a heavy chain by disulfide bonding through constant regions. The light and heavy chain variable regions of a Fab region provide for an Fv region that participates in antigen binding.

The antibody variable region can be present in a recombinant derivative. Examples of recombinant derivatives include single-chain antibodies, diabody, triabody, tetrabody, and miniantibody. (Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004.)

The antigen binding protein can contain one or more variable regions recognizing the same or different epitopes. (Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004.)

II. Generation of Antigen Binding Protein Directed to an Identified Target Region

Different antigen binding proteins directed to the mAb 1G3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1 target region, or mAb 13G11.BF3 target region can be generated starting with the respective monoclonal antibody. Alternatively, the epitope recognized by a binding protein can be used to select additional binding proteins.

The mAb 2H2.BE11 target region appears to be located at approximately amino acids 76-357 of ORF0657n. A polypeptide containing amino acids 76-357 of ORF0657n, or a full-length ORF0657n, can be used as a target antigen to select for antibodies. The target region of the generated antibodies can be determined.

A variety of techniques are available to select for a protein recognizing an antigen. Examples of such techniques include use of phage display technology and hybridoma production. Human antibodies can be produced using chimeric mice such as a XenoMouse or Trans-Chromo mouse.

(E.g., Azzazy *et al.*, *Clinical Biochemistry* 35:425-445, 2002, Berger *et al.*, *Am. J. Med. Sci.* 324(1):14-40, 2002.)

The monoclonal antibodies mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 contain variable regions recognizing ORF0675n. Additional binding proteins recognizing ORF0657n can be produced based on antibody variable regions. Additional binding proteins can, for example, be produced by modifying an existing monoclonal antibody and by using variable region sequence information. Protein construction and sequence manipulation can be performed using recombinant nucleic acid techniques.

The monoclonal antibodies mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, and mAb 13G11.BF3 are murine antibodies. For human therapeutic applications, preferred binding proteins based on such mAb's are designed to reduce the potential generation of human anti-mouse antibodies recognizing the murine regions.

The potential generation of human anti-mouse antibodies can be reduced using techniques such as murine antibody humanization, de-immunization, and chimeric antibody production. (See, for example, O'Brien *et al.*, Humanization of Monoclonal Antibodies by CDR Grafting, p 81-100, From *Methods in Molecular Biology* Vol. 207: Recombinant antibodies for Cancer Therapy: Methods and Protocols (Eds. Welschhof and Krauss) Humana Press, Totowa, New Jersey, 2003; Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004; Gonzales *et al.*, *Tumor Biol.* 26:31-43, 2005, Presta, *Advanced Drug Delivery Reviews* 58:640-656, 2006, Tsurushita *et al.*, *Methods* 36:69-83, 2005, Roque *et al.*, *Biotechnol. Prog.* 20:639-654, 2004.)

Murine antibodies can be humanized using techniques such as grafting complementary determining regions into a framework region or resurfacing. Resurfacing (also known as veneering) involves modifying a variable region so the surface exposed regions are humanized.

Grafting complementary determining regions involves taking such regions or a portion of such regions from, for example, a murine source and inserting the regions into a human variable region framework. The human framework used for grafting can be selected based on sequence homology to the variable region (e.g., murine) from which the region was obtained. Essential framework residues associated with grafted complementary determining regions should also be provided in the new framework.

De-immunization involves altering potential linear T-cell epitopes present in the antibody. The epitopes can be identified based on a bioinformatics scan of known human HLA class I and/or class II epitopes. (Presta, *Advanced Drug Delivery Reviews* 58:640-656, 2006.)

A chimeric antibody contains a human constant region along with a variable region from a different organism, such as a mouse. The human constant region provides an Fc region.

Additional examples of alterations include providing a variable region in, for example, a single chain antibody, a diabody, a triabody, a tetrabody, and a miniantibody. (Kipriyanov *et al.*, *Molecular Biotechnology* 26:39-60, 2004.) The antigen binding protein can contain one or more variable

regions recognizing the same or different epitopes. (*Id.*) Additional embodiments of the present invention are directed to a single chain antibody, a diabody, a triabody, a tetrabody, or a miniantibody directed to the mAb 1G3.BD4, mAb 2H2.BE11, mAb 13C7.BC1, or mAb 13G11.BF3 binding site.

5 III. Binding Protein Directed to the mAb 2H2.BE11 Target Region

As described in the Examples provided below, the mAb 2H2.BE11 target region was further characterized and the amino acids sequence of the variable regions was determined. The identified target region and the sequence information facilitate obtaining different binding proteins directed to the mAb 2H2.BE11 target region.

10 In an embodiment of the present invention, the binding protein binds to a polypeptide consisting of amino acids 76-357 of SEQ ID NO: 1. Preferably, the binding protein is either a human antibody, a humanized antibody, a de-immunized antibody, or chimeric antibody. Preferred antibodies are isolated antibodies and monoclonal antibodies.

The amino acids sequences of the mAb 2H2.BE11 variable regions are provided by SEQ
15 ID NO: 20 (V_h) and SEQ ID NO: 21 (V_l). The complementary determining regions (CDR's) within V_h were identified at amino acids 36-45, 50-65, and 98-107. The CDR's within V_l were identified at amino acids 24-33, 49-55, and 88-96 of SEQ ID NO: 21.

In different embodiments directed to a V_h region, the binding protein binds the mAb 2H2.BE11 target region and comprises, consists, or consists essentially of: a first V_h CDR comprising,
20 consisting, or consisting essentially of amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid; a second V_h CDR comprising, consisting, or consisting essentially of amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acid; and a third V_h CDR comprising, consisting, or consisting essentially of amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.

25 In different embodiments directed to a V_l region, the binding protein binds the mAb 2H2.BE11 target region and comprises, consists, or consists essentially of a first V_l CDR comprising, consisting, or consisting essentially of amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid; a second V_l CDR comprising, consisting, or consisting essentially of amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by
30 one amino acid; and a third V_l CDR comprising, consisting, or consisting essentially of amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.

Reference to "consisting essentially of" with respect to a variable region, CDR region, or antibody sequence, indicates the possible presence of one or more additional amino acids, where such amino acids do not significantly decrease binding to the target.

35 An amino acid difference can be an amino acid deletion, insertion, or substitution. In substituting amino acids to maintain activity, the substituted amino acids should have one or more similar properties such as approximately the same charge, size, polarity and/or hydrophobicity.

Preferably, an amino acid substitution is a conservative substitution. A conservative substitution replaces an amino acid with another amino acid having similar properties. Table 1 provides a list of groups of amino acids, where one member of the group is a conservative substitution for another member.

5

Table 1 : Conservative Substitutions

Ala, Val, Ile, Leu, Met
Ser, Thr,
Tyr, Trp
Asn, Gln
Asp, Glu
Lys, Arg, His

In additional embodiments the V_h region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and/or the V_l region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

10

In different embodiments focusing on an antibody, the antibody comprises, consists, or consists essentially of: (a) a heavy chain comprising a V_h region as described in this Section III, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄, and (b) a light chain comprising a V_l region as described above in this section III, and a human kappa C_L or human lambda C_L. In further embodiments: the antibody comprises, consists, or consists essentially of: (a) a heavy chain comprising a V_h region as described in this Section III, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁ or IgG₂ and (b) a light chain comprising a V_l region as described above in this Section III, and a human kappa C_L; and the heavy chain consists essentially of the amino acid sequence of SEQ ID NO: 22 and/or the light chain consists essentially of the amino acid sequence of SEQ ID NO: 23.

15

20

In additional embodiments the antigen-binding protein described herein has V_h and V_l regions providing an affinity K_D at least about 100 nM, preferably at least about 30 nM to the target antigen. Binding to the target antigen can be determined as described in Example 11, using an ORF0657n fragment from amino acids 42-486

25

Preferred binding proteins for the different embodiments are an antibody. More preferably the antibody is isolated or a monoclonal antibody.

IV. Protein Production

Antigen binding protein are preferably produced using recombinant nucleic acid techniques or through the use of a hybridoma. Recombinant nucleic acid techniques involve constructing

a nucleic acid template for protein synthesis. Hybridoma techniques involve using an immortalized cell line to produce the antigen binding protein. Suitable recombinant nucleic acid and hybridoma techniques are well known in the art. (See for example, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 2005, Harlow *et al.*, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988.)

5 Recombinant nucleic acid encoding an antigen binding protein can be expressed in a host cell that in effect serves as a factory for the encoded protein. The recombinant nucleic acid can provide a recombinant gene encoding the antigen binding protein that exists autonomously from a host cell genome or as part of the host cell genome.

10 A recombinant gene contains nucleic acid encoding a protein along with regulatory elements for protein expression. Generally, the regulatory elements that are present in a recombinant gene include a transcriptional promoter, a ribosome binding site, a terminator, and an optionally present operator. A preferred element for processing in eukaryotic cells is a polyadenylation signal. Antibody associated introns may also be present. Examples of expression cassettes for antibody or antibody fragment production are well known in art. (*E.g.*, Persic *et al.*, *Gene* 187:9-18, 1997, Boel *et al.*, *J. Immunol. Methods* 239:153-166, 2000, Liang *et al.*, *J. Immunol. Methods* 247:119-130, 2001, Tsurushita *et al.*, *Methods* 36:69-83, 2005.)

Due to the degeneracy of the genetic code, a large number of different encoding nucleic acid sequences can be used to code for a particular protein. The degeneracy of the genetic code arises because almost all amino acids are encoded by different combinations of nucleotide triplets or "codons".

20 Amino acids are encoded by codons as follows:

A=Ala=Alanine: codons GCA, GCC, GCG, GCU

C=Cys=Cysteine: codons UGC, UGU

D=Asp=Aspartic acid: codons GAC, GAU

E=Glu=Glutamic acid: codons GAA, GAG

25 F=Phe=Phenylalanine: codons UUC, UUU

G=Gly=Glycine: codons GGA, GGC, GGG, GGU

H=His=Histidine: codons CAC, CAU

I=Ile=Isoleucine: codons AUA, AUC, AUU

K=Lys=Lysine: codons AAA, AAG

30 L=Leu=Leucine: codons UUA, UUG, CUA, CUC, CUG, CUU

M=Met=Methionine: codon AUG

N=Asn=Asparagine: codons AAC, AAU

P=Pro=Proline: codons CCA, CCC, CCG, CCU

Q=Gln=Glutamine: codons CAA, CAG

35 R=Arg=Arginine: codons AGA, AGG, CGA, CGC, CGG, CGU

S=Ser=Serine: codons AGC, AGU, UCA, UCC, UCG, UCU

T=Thr=Threonine: codons ACA, ACC, ACG, ACU

V=Val=Valine: codons GUA, GUC, GUG, GUU

W=Trp=Tryptophan: codon UGG

Y=Tyr=Tyrosine: codons UAC, UAU

Expression of a recombinant gene in a cell is facilitated using an expression vector.

- 5 Preferably, the expression vector, in addition to a recombinant gene, also contains an origin of replication for autonomous replication in a host cell, a selectable marker, a limited number of useful restriction enzyme sites, and a potential for high copy number. Examples of expression vectors for antibody and antibody fragment production are well known in art. (*E.g.*, Persic *et al.*, *Gene* 187:9-18, 1997, Boel *et al.*, *J. Immunol. Methods* 239:153-166, 2000, Liang *et al.*, *J. Immunol. Methods* 247:119-130, 2001, Tsurushita *et al.*, *Methods* 36:69-83, 2005.)

10 If desired, nucleic acid encoding an antibody may be integrated into the host chromosome using techniques well known in the art. (*E.g.*, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 2005, Marks *et al.*, International Application Number WO 95/17516, International Publication Date June 29, 1995.)

- 15 A variety of different cell lines can be used for recombinant antigen binding protein expression, including those from prokaryotic organisms (*e.g.*, *E. coli*, *Bacillus sp.*, and *Streptomyces sp.* (or streptomycete) and from eukaryotic (*e.g.*, yeast, Baculovirus, and mammalian). (Breitling *et al.*, *Recombinant Antibodies*, John Wiley & Sons, Inc. and Spektrum Akademischer Verlag, 1999, Kiprianov *et al.*, *Molecular Biotechnology* 26:39-60, 2004, Tsurushita *et al.*, *Methods* 36:69-83, 2005.)

- 20 Preferred hosts for recombinant antigen binding protein expression provide for mammalian post translational modifications. Post translational modifications chemical modification such as glycosylation and disulfide bond formation. Another type of post translational modification is signal peptide cleavage.

- 25 Proper glycosylation can be important for antibody function. (Yoo *et al.*, *Journal of Immunological Methods* 261:1-20, 2002, Li *et al.*, *Nature Biotechnology* 24(2):210-215, 2006.) Naturally occurring antibodies contain at least one N-linked carbohydrate attached to a heavy chain. (Yoo *et al.*, *Journal of Immunological Methods* 261:1-20, 2002.) Additional N-linked carbohydrates and O-linked carbohydrates may be present and may be important for antibody function. (*Id.*)

- 30 Different types of host cells can be used to provide for efficient post-translational modifications including mammalian host cells and non-mammalian cells. Examples of mammalian host cells include but are not limited to Chinese hamster ovary (Cho), HeLa, C6, PC12, Human Embryonic Kidney (HEK293) and myeloma cells. (Yoo *et al.*, *Journal of Immunological Methods* 261:1-20, 2002, Persic *et al.*, *Gene* 187:9-18, 1997.) Non-mammalian cells can be modified to replicate human glycosylation. (Li *et al.*, *Nature Biotechnology* 24(2):210-215, 2006.) Glycoengineered *Pichia pastoris* is an example of such a modified non-mammalian cell. (Li *et al.*, *Nature Biotechnology* 24(2):210-215, 2006.)
- 35

Preferred recombinant genes comprise a nucleotide sequence encoding an antibody variable region that binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region. A particular recombinant gene can encode for a protein containing one variable region or both a V_H and V_L region.

- 5 The recombinant gene can also encode for antibody constant regions and hinge region. If desired, an antibody can be produced using a combination of recombinant genes, where one gene encodes for a light chain and the second gene encodes for a heavy chain.

Different embodiments are provided by the nucleic acid encoding a protein described in Section II or III *supra*. Examples of such embodiments are provided below.

- 10 In an embodiment directed to a V_H encoding region, the nucleotide sequence encodes a variable region comprising, consisting, or consisting essentially of: a first V_H CDR comprising, consisting, or consisting essentially of amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid; a second V_H CDR comprising, consisting, or consisting essentially of amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acid; and a third V_H CDR comprising, consisting, or consisting essentially of amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.

- In an embodiment directed to a V_L encoding region, the nucleotide sequence encodes a variable region comprising, consisting, or consisting essentially of a first V_L CDR comprising, consisting, or consisting essentially of amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid; a second V_L CDR comprising, consisting, or consisting essentially of amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and a third V_L CDR comprising, consisting, or consisting essentially of amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.

- 20 In additional embodiments: the V_H region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and the V_L region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

- In different embodiments focusing on an antibody heavy and/or light chain, the recombinant gene encodes either or both a protein comprising, consisting, or consisting essentially of: (a) a heavy chain comprising a V_H region as provided in Section III *supra*, a human hinge, CH₁, CH₂, and CH₃ from an IgG1, IgG2, IgG3 or IgG4 subtype or (b) a light chain comprising a V_L region as provided in Section III *supra*, and a human kappa C_L or lambda C_L. In a further embodiment the heavy chain consists essentially of the amino acid sequence of SEQ ID NO: 22; and the light chain consists essentially of the amino acid sequence of SEQ ID NO: 23.

35 V. Applications of Antigen Binding Proteins.

Antigens containing certain ORF0657n regions can be used to provide a protective immune response against *S. aureus* infection. (Anderson *et al.*, International Publication No. WO

2005/009379, International Publication Date February 3, 2005.) An antigen binding protein recognizing an ORF0657n target region can be used to facilitate the production, characterization, or study of ORF0657n antigens and vaccines. Antigen binding protein recognizing appropriate epitopes can also have therapeutic applications.

5 Examples of different uses in the production, characterization, or study of ORF0657n related antigens and vaccines include:

- 1) Identifying the presence of an ORF0657n antigen, for example, by Western blot;
- 2) Identifying the presence of an ORF0657n antigen on a cell surface, for example, by
10 flow cytometry. This is useful, for example, in determining expression on multiple strains of *S. aureus* as well as confirmation of knock-out mutants;
- 3) Passive protection experiments. The antibodies can be used in a lethal model to determine if a specific area of the ORF0657n protein confers protection;
- 4) An immunoassay. The assay can be used to monitor antigen quality, product
15 production and stability;
- 5) As a control in mouse potency assays to monitor immunogenicity of an antigen vaccine product; and
- 6) Serology assays can utilize a monoclonal antibody in a competitive format to identify an immune response to ORF0657n derived antigen vaccinated patients.

20 Techniques for using antigen binding proteins, such as monoclonal antibodies, in the production, characterization, or study of a target protein are well known in the art. (See, for example, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 2005, Harlow *et al.*, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988, Harlow *et al.*, *Using Antibodies*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., Cold Spring Harbor Laboratory Press, 1999, Lipman *et al.*, *ILAR Journal* 46:258-268, 2005.)

25 In an embodiment of the present invention, the presence of an ORF0657n antigen in a solution, bound to a microsphere or on a cell is determined using an antigen binding protein. The ability of the binding protein to bind to a protein present in the solution or cell can be determined using different techniques such as a Western blot, enzyme-linked immunosorbent assay (ELISA), flow cytometry, and Luminex immunoassay.

30 VI. Treatment

35 Therapeutic and prophylactic treatment can be performed on a patient using an antigen binding protein binding to an appropriate target region. Therapeutic treatment is performed on those persons infected with *S. aureus*. Prophylactic treatment can be performed on the general population or a subset of the general population. A preferred subset of the general population are those persons at an increased risk of *S. aureus* infection.

A "patient" refers to a mammal capable of being infected with *S. aureus*. Preferably, the patient is a human. However, other types of mammals such as cows, pigs, sheep, goats, rabbits, horses, dogs, cats, monkeys, rats, and mice, can be infected with *S. aureus*. Treatment of non-human patients is useful in protecting pets and livestock, and in evaluating the efficacy of a particular treatment.

5 Persons with an increased risk of *S. aureus* infection include health care workers; hospital patients; patients with a weakened immune system; patients undergoing surgery; patients receiving foreign body implants, such as a catheter or a vascular device; patients facing therapy leading to a weakened immunity; and persons in professions having an increased risk of burn or wound injury. (*The Staphylococci in Human Disease*, Crossley and Archer (ed.), Churchill Livingstone Inc. 1997.)

10 In an embodiment, a patient is administered an antigen binding protein in conjunction with surgery or a foreign body implant. Reference to "surgery or a foreign body implant" includes surgery with or without providing a foreign implant, and providing a foreign implant with or without surgery. The timing of administration can be designed to achieve prophylactic treatment and/or therapeutic treatment. Administration is preferably started around the same time as surgery or
15 implantation.

Guidelines for pharmaceutical administration in general are provided in, for example, *Remington's Pharmaceutical Sciences 20th Edition*, Ed. Gennaro, Mack Publishing, 2000; and *Modern Pharmaceutics 2nd Edition*, Eds. Banker and Rhodes, Marcel Dekker, Inc., 1990.

20 Pharmaceutically acceptable carriers facilitate storage or administration of an antigen binding protein. Substances used to stabilize protein solution formulations include carbohydrates, amino acids, and buffering salts. (Middaugh *et al.*, *Handbook of Experimental Pharmacology 137:33-58*, 1999.)

Antigen binding proteins can be administered by different routes such as intravenous, subcutaneous, intramuscular, or mucosal. Subcutaneous and intramuscular administration can be performed using, for example, needles or jet-injectors. Mucosal delivery, such as nasal delivery, can
25 involve using enhancers or mucoadhesives to produce a longer retention time at adsorption sites. (Middaugh *et al.*, *Handbook of Experimental Pharmacology 137:33-58*, 1999.)

Suitable dosing regimens are preferably determined taking into account factors well known in the art including age, weight, sex and medical condition of the patient; the route of administration; the desired effect; and the particular compound employed. It is expected that an effective
30 dose range should be about 0.1 mg/kg to 20 mg/kg, or 0.5 mg/kg to 5 mg/kg. The dosing frequency can vary depending upon the effectiveness and stability of the compound. Examples of dosing frequencies include biweekly, weekly, monthly and bimonthly.

VII. Examples

35 Examples are provided below further illustrating different features of the present invention. The examples also illustrate useful methodology for practicing the invention. These examples do not limit the claimed invention.

Example 1: Generation of Monoclonal Antibodies to ORF0657n

Monoclonal antibodies directed to ORF0657n (SEQ ID NO: 1) were generated using ORF0657n-C/e (SEQ ID NO: 2) or ORF0657n-H/y (SEQ ID NO: 3) as an antigen. The antibodies were identified and characterized by ELISA and flow cytometry.

Mice and Immunizations: Female BALB/c mice, 4-5 weeks old, were purchased from Taconic (Germantown, N. Y.). Mice were immunized intramuscularly (i.m.) on days 0, 7, and 21, with 20 µg of *E. coli* produced ORF0657n-C/e antigen or Yeast expressed ORF0657n-H/y antigen, formulated on aluminum hydroxyphosphate adjuvant. (Anderson *et al.*, International Publication No. WO 2005/009379, International Publication Date February 3, 2005.) A final intravenous injection (i.v.) of 20 µg of protein in phosphate buffered saline (PBS) was given to mice three days prior to the fusion. Mice were sacrificed and the spleens removed for cell fusion.

Mab Production: Lymphocytes prepared from spleens were fused with the mouse myeloma partner SP2/0-Ag14 (ATCC 1581) by polyethylene glycol 1500 (Boehringer Mannheim) at a ratio of 3:1. The fusions were plated into 96-well flat-bottomed microtiter plates in Dulbecco's Modification of Eagle's Medium, high glucose, pyruvate (DMEM) containing 20% fetal bovine serum, hypoxanthine (10^{-4} M), thymidine (10^{-5} M), Aminopterin (4×10^{-7} M) was added 24 hours later. Supernatants from growing hybridomas were screened by ELISA for reactivity to ORF0657n as described below. Positive wells were cloned by limiting dilution and retested for ELISA reactivity. Monoclonal antibodies were classified with an antibody-isotyping kit (Roche Diagnostics Corporation, Indianapolis, IN).

ELISA: Costar medium binding microtiter plates were coated overnight at 2-8°C with 50 nanograms per well of *E. coli* expressed SEQ ID NO: 2 in PBS. The plate was washed three times with PBS, 0.05% Tween20 and blocked with 1% Bovine serum albumin, PBS, 0.05% Tween20 (assay diluent) for at least 1 hour. The plate was washed as before and supernatants from the fusion wells or cloned hybridomas were added and allowed to incubate for 2 hours at room temperature. The plate was washed as before and a Goat anti-mouse IgG (H+L)-HRP conjugate (Zymed) (1:8000 in assay diluent) added and allowed to incubate for 1 hour at room temperature. Assay plates were developed with TMB substrate, the reaction stopped with 2.0 N H₂SO₄ and read in a plate reader at OD 450 nm. Wells were considered positive that had an optical density at 450 nm of >1.0.

Flow Cytometry: Prepared glycerol stocks of *S. aureus* passaged under iron-starved conditions (in RPMI) were used to evaluate mAb for ORF0657n binding. Frozen glycerol stock cells were thawed and resuspended in PBS; 1% bovine serum albumin; 0.1% sodium azide, 0.2% Pig IgG (Sigma) (PAAG) to a concentration of 5×10^7 CFU/50 µl. A 50 µl aliquot of the cells were placed in a 1.5 ml Eppendorf tube per reaction. Fifty microliters of the hybridoma culture were added to each reaction tube and incubated for 1 hour at room temperature. The cells were washed by adding 1 mL of phosphate buffered saline; 1% bovine serum albumin; 0.1% sodium azide (PAA) to the tube. The cells

were pelleted by centrifugation (5500 rpm, 5 minutes). The supernatant was removed and the cells were mixed with 100 μ L of secondary antibody (FITC-labeled goat anti-mouse Ig (BD Pharmingen) diluted 1:100 in PAAG). Incubation was for 1 hour at room temperature in the dark. After incubation, 1 mL PAA was added to the reaction mixture, the cells were pelleted (5500 rpm, 5 minutes) and supernatant removed. The pellets were resuspended in 1 mL of PBS and transferred to 12 x 75 mm tubes for FAC analysis.

Tubes were run on a BD-FACSCalibur flow cytometer instrument gated for bacterial cells and measuring the amount of FITC associated with the cells. A standard antibody with known binding to the surface of *S. aureus* was run in every assay. A negative control was run as cells and the secondary conjugate alone. Hybridoma wells were considered positive if the geometric mean value was greater than 30.

Two separate fusions resulted in a panel of twelve monoclonal antibodies (mAb). All of the mAbs were reactive in ELISA (Table 2). Ten of the twelve mAbs bound to the surface of bacteria as evidenced by flow cytometry. All of the mAbs were positive by Western Blot analysis with the wild type protein.

Table 2

mAbs/cell lines Fusion #1	mAbs/cell lines Fusion #2
1) 2H2.B8 IgG1	
2) 8H6.E11.H3 IgG2a*	
3) 7H2.C11 IgG1*	
	4) 2E12.A8 IgG1
	5) 8A8.B4 IgG1
	6) 3G11.D5 IgG1
	7) 13G11.C11 IgG1
	8) 13C7.D12 IgG1
	9) 1G3.B3 IgG1
	10) 9H3.E4 IgG1
	11) 3B7.G8 IgG1
	12) 3G12.A4 IgG1

* Not reactive in flow cytometry. Fusion #1 was generated from *E. coli* produced ORF0657n-C/e antigen. Fusion #2 was generated with Yeast expressed ORF0657n-H/y antigen.

Example 2: Class Switching mAbs

All of the mAbs isolated that bound to the native antigen were of the IgG1 isotype. These antibodies were class switched to an IgG2b isotype by selecting for shift variants (Spira *et al*, *J. of Immunological Methods*, 74:307-315, 1985). A suitable immunoassay was developed using an IgG2b conjugate and the cell line was plated at a high density. Somatic cell mutations were selected, enriched

and then cloned. The binding site of the switched mAb remained identical to the original mAb, but switching to an IgG2b subtype gave a more favorable isotype (initiating the complement cascade) in the passive protection studies.

5 Table 3 Class Switched mAbs

IgG1 isotype	IgG2b isotype
2H2.B8	2H2.BE11
2E12.A8	2E12.BG1
8A8.B4	8A8.BF9
3G11.D5	3G11.BE5
13G11.C11	13G11.BF3
13C7.D12	13C7.BC1
1G3.B3	1G3.BD4
9H3.E4	9H3.BE4

Example 3: Binding Inhibition Studies with Native Antigen

Purified antibodies were labeled with Alexafluor-488 using a mAb labeling kit
 10 (Molecular Probes) according to the manufacturer's instructions. The amount of mAb that would just saturate the surface of RPMI-grown bacterial cells was determined for both the labeled and unlabeled mAbs. Each of the mAbs in Table 3 (1st column) were used labeled and unlabeled.

The inhibition assay was performed by first incubating 5×10^7 cells with the unlabeled mAb at a concentration that would saturate the surface of the cells. This reaction was incubated at room
 15 temperature for 1 hour. After this incubation, the reactions were washed with 1 ml of PAA and spun at 6,000 RPM for 5 minutes in a microcentrifuge (Hermle). The supernatant was removed down to ~50 ul and the cells were resuspended in 100 ul of PAAG containing the amount of directly labeled mAb that would just saturate the surface of the cells. After this incubation, the reactions were washed with 1 ml of PAA and spun at 6,000 RPM for 5 minutes in a microcentrifuge (Hermle). The supernatant was
 20 removed down to ~50 ul and the cells were resuspended in 1 ml of PBS and transferred to 12 x 75 mm tubes for FAC analysis. As controls, separate reactions with the unlabeled mAb were measured with a secondary Alexafluor-488 conjugated goat anti-mouse IgG (H+L) (Molecular probes, 1:400 in PAAG) to determine that this mAb was bound to the surface. A positive control was also performed that had only the labeled mAb with the cells. If the unlabeled mAb bound to the same epitope as the labeled mAb then
 25 there would be no or low fluorescent reactivity associated with the cells. If the unlabeled mAb bound to a different epitope than the labeled mAb then the level of reactivity associated with the surface would be equivalent to the labeled mAb only control cells.

The panel of monoclonal antibodies fell into four reactive groups by inhibition studies:

Table 4

Group I	Group II	Group III	Group IV
2H2.B8	9H3.E4	13G11.C11	2E12.A8
8A8.B4	1G3.B3		13C7.D12
	3G11.D5		

5

Example 4: Binding Studies with Denatured Antigen and Altered Antigens

ORF0657n altered proteins were used to further characterize binding. Nucleic acid encoding ORF0657n was initially cloned into the expression vector pET-28a (Novagen) and expressed in *E. coli* with a C-terminal 6X his tag (SEQ ID NO: 2). The expression vector with the cloned gene was subjected to mutagenesis using Stratagene's QuikChange XL Site-Directed Mutagenesis Kit following the manufacturer's instructions. The gene was mutated with specific sequential amino acid changes. The resulting plasmid was transformed into Stratagene's XL10-Gold competent cells following the manufacturer's protocol. Plasmids were isolated from transformants using Qiagen's QIAprep Spin Miniprep Kit. Transformants were screened by sequencing using ABI's 310 DNA Sequencer. Plasmid from the transformant exhibiting the greatest number of base changes was transformed into the expression host HMS174(DE3) (Novagen). Transformants were expressed following Novagen's instructions.

10

15

20

Different ORF0657n altered proteins were used to determine the diversity of the ORF0657n mAbs (SEQ IDs 4-19). These proteins were screened with the 10 different mAbs in dot blots using standard procedures. Positive/negatives were confirmed by Western blots using standard procedures. By this approach antibodies were grouped according to their binding profile. Seven of the antibodies resolved to three groups; the three remaining antibodies (2H2.B8, 8A8.E11.H3 and 13G11.C11) had profiles that were similar but not identical to each other (Table 5).

TABLE 5: Binding of ORF0657n specific mAbs to ORF0657n mutant proteins detected by Western blot

	Group III		Group II			Group IV		Group I		
SEQ ID NO:	3G11.C11	3G12.A4	3B7.G8	1G3.B3	9H3.E4	2E12.A8	13C7.D12	2H2.B8	8A8.E11.H3	13G11.C11
1	+	+	+	+	+	+	+	+	+	+
2	+	+	+	+	+	+	+	+	+	+
3	+	+	+	+	+	+	+	+	+	+

TABLE 5: Binding of ORF0657n specific mAbs to ORF0657n mutant proteins detected by Western blot

SEQ ID NO:	Group III		Group II			Group IV		Group I		
	3G11.C11	3G12.A4	3B7.G8	1G3.B3	9H3.E4	2E12.A8	13C7.D12	2H2.B8	8A8.E11.H3	13G11.C11
4	+	+	+	+	+	+	+	+	+	+
5	+	+	+	+	+	+	+	+	+	+
6	+	+	+	+	+	-	-	+	+	+
7	+	+	+	+	+	-	-	+	+	+
8	+	+	+	+	+	-	-	+	+	+
9	+	+	+	+	+	-	-	-	+	+
10	+	+	+	+	+	-	-	-	+	+
11	-	-	W	W	W	-	-	-	-	W
12	-	-	W	W	W	-	-	-	-	W
13	-	-	-	-	-	-	-	-	-	-
14	+	+	+	+	+	+	+	+	+	+
15	+	+	+	+	+	+	+	+	+	+
16	+	+	+	+	+	+	+	-	+	+
17	+	+	+	+	+	+	+	-	+	+
18	+	+	+	+	+	+	+	-	+	-
19	+	+	+	+	+	W	W	-	-	+

+, Antibody bound to protein in a Western; -, Antibody did not bind to protein by Western; W, Weak binding of antibody to protein detected by Western. Antibodies were grouped according to hybridization profile. A dotted line is used where similar, but not identical profiles were obtained.

Example 5: BIAcore Studies

In BIAcore studies the mAbs were examined by "footprint analysis" using purified ORF0657n-H/y as the antigen. Pair-wise binding experiments were conducted using real-time biomolecular interaction analysis via BIACORE[®]. BIACORE[®] incorporates microfluidics technology and surface plasmon resonance (SPR) to detect changes in mass by monitoring changes in the refractive index of a polarized light aimed directly at the surface of a carboxyl methyl dextran coated (CM5) sensor chip. The changes in response, measured in Response Units, can be correlated to the amount of bound analyte (*i.e.* antigen or antibody).

An anti-staphylococcal antibody (mAb 13C7.D12) was covalently bound (immobilized) on the surface of the CM5 sensor chip. The immobilized Ab was exposed first to the ORF0657n protein and subsequently to a pair of antibodies in a matrix format. After each cycle of ORF0657n protein + antibody pair, the surface of the sensor chip was regenerated back to the immobilized mAb 13C7.D12

using 20 mM HCl. Eight antibodies were tested against the ORF0657n protein in a matrix format so that all combinations of each antibody pair could be analyzed. The matrix design for mAb pairs used in this experiment is summarized in Table 6.

Table 6. Summary of Antibodies Tested in 8x8 Matrix

Cycle #	First Antibody	Second Antibody			
		Flow Cell 1	Flow Cell 2	Flow Cell 3	Flow Cell 4
1	N/A Immobilization	13C7.D12	13C7.D12	13C7.D12	13C7.D12
2	2H2.B8	2H2.B8	13C7.D12	8A8.B4	9H3.E4
3	2H2.B8	13G11.C11	2E12.A8	1G3.B3	3G11.D5
4	13C7.D12	2H2.B8	13C7.D12	8A8.B4	9H3.E4
5	13C7.D12	13G11.C11	2E12.A8	1G3.B3	3G11.D5
6	8A8.B4	2H2.B8	13C7.D12	8A8.B4	9H3.E4
7	8A8.B4	13G11.C11	2E12.A8	1G3.B3	3G11.D5
8	9H3.E4	2H2.B8	13C7.D12	8A8.B4	9H3.E4
9	9H3.E4	13G11.C11	2E12.A8	1G3.B3	3G11.D5
10	13G11.C11	2H2.B8	13C7.D12	8A8.B4	9H3.E4
11	13G11.C11	13G11.C11	2E12.A8	1G3.B3	3G11.D5
12	2E12.A8	2H2.B8	13C7.D12	8A8.B4	9H3.E4
13	2E12.A8	13G11.C11	2E12.A8	1G3.B3	3G11.D5
14	1G3.B3	2H2.B8	13C7.D12	8A8.B4	9H3.E4
15	1G3.B3	13G11.C11	2E12.A8	1G3.B3	3G11.D5
16	3G11.D5	2H2.B8	13C7.D12	8A8.B4	9H3.E4
17	3G11.D5	13G11.C11	2E12.A8	1G3.B3	3G11.D5

5

To normalize for the amount of antigen initially bound (captured) in each run, the following ratio for each test antibody/antigen complex is calculated:

$$= \frac{\text{Test Antibody Response Units} * 1000}{\text{ORF0657n protein Response Units}} \quad \text{or} \quad \frac{\text{mRU}_{\text{Ab}}}{\text{RU}_{\text{Ag}}}$$

10

The percentage of available epitope remaining for each antibody can be calculated for the mapping pair as follows:

$$= \frac{(\text{mRU}_{\text{Ab}} (\text{when } 2^{\text{nd}} \text{ Ab}) / \text{RU}_{\text{Ag}}) * 100}{(\text{mRU}_{\text{Ab}} (\text{when } 1^{\text{st}} \text{ Ab}) / \text{RU}_{\text{Ag}})} \quad \text{or} \quad \begin{array}{l} \% \text{ Remaining} \\ \text{(calculated for each Ab)} \end{array}$$

Figure 2 illustrates matrix resulting outlining the reactivities of the monoclonal antibodies in a pair-wise binding study. The panel of monoclonal antibodies fell into three reactive areas by the BIAcore® method (See Table 7).

5

Table 7

Group I	Group II	Group III
2H2.B8	13G11.C11	13C7.D12
8A8.B4	3G11.D5	2E12.A8
9H3.E4		
1G3.B3		

Example 6: Protection Studies with Passive Immunization in a Murine Sepsis Model

10

The monoclonal antibodies mAb 2H2.BE11 and mAb 13C7.BC1 were tested for their ability to provide protection against *S. aureus* infection. These antibodies recognize different epitopes on the ORF0657n protein. Controls included an isotype matched mAb and PBS-only.

15

The mAbs or PBS were administered intraperitoneally (i.p.) 20 hours prior to bacterial challenge. Mice were then challenged with a LD₈₀₋₉₀ dose of *S. aureus* Becker i.v. and monitored for survival. Each experiment was repeated three times with groups of 10 or 20 mice and was monitored for 10 days. The half life for the monoclonal antibodies in uninfected BALB/c mice is approximately eight days. A dose of 0.5 mg was found to be optimal. The results of experiments with the two monoclonal antibodies are presented in Figures 3A-C, 4A, 4B, and 5A-C.

20

Whereas the mAb 13C7.BC1 significantly improved survival at day 10 compared to the controls in one experiment, in the other 2 repetitions the overall survival rate was similar to that of the controls (Figures 3A-3C). However, compared to controls, there was delay in the time to death of the mAb 13C7.BC1 treated mice within this 10 day period. A similar trend in delay of time to death of the mAb 2H2.BE11 treated mice was also noted in two of the three experiments (Figures 5A-5C).

25

The effect of mAb 13C7.BC1 was also examined using a recent *S. aureus* clinical isolate UK58 (Figures 4A and 4B). This strain was minimally passaged from an abscess site in a patient. In two independent experiments, the results show a delay in time to death with the UK58 challenge.

30

Antibody persistence studies cannot be evaluated in the LD₈₀₋₉₀ model due to the rapid rate of death. Therefore, a sub-lethal challenge model was run. In the sub-lethal model the challenge dose used is 10% of that used for the LD₈₀₋₉₀ model. The sub-lethal challenge model was monitored over a four day period. Groups of 22 mice received 0.5 mg doses of either mAb 13C7.BC1 or isotype control mAb (6G6) 20 hours prior to i.v. bacterial challenge with 5×10^7 CFU of *S. aureus* Becker. Two animals from each group were sacrificed just prior to challenge (T=0) to determine the mAb levels in the

serum at the time of challenge. At 2, 24, 48, 72 and 96 hours post challenge, four mice from each group were sacrificed and serum mAb levels determined.

From this sub-lethal challenge experiment, the half life of mAb 13C7.BC1 in *S. aureus*-infected mice was estimated to be approximately one-day. In contrast, the half life of the isotype control mAb was estimated to be greater than four days (data not shown). These data point to a specific reduction of mAb 13C7.BC1 in *S. aureus* challenged mice, which appears to be exhausted well before the ten day period monitored in the lethal model.

In six of the eight experiments illustrated in Figures 3A-C, 4A, 4B, and 5A-C, improved survival was observed through approximately three days for the groups receiving the mAb administration. These results provide an indication that such mAbs have a positive effect on the survival rate of *S. aureus* challenged mice.

Example 7: Protection Studies with Passive Immunization in a Murine Indwelling Catheter Model

A murine indwelling catheter model was used with mAb 2H2.BE11. The *S. aureus* strain used in this model was the clinical isolate MCL8538. This strain was selected as lower inocula could be administered while still getting reproducible colonization of catheters compared to *S. aureus* Becker, the strain used in the murine sepsis model.

ICR mice had catheters (PE50 silicone rubber) surgically implanted into the jugular vein, held in place with sutures, and exiting with a port on the dorsal midline of the mouse. Mice were rested 9-11 days post surgery. At 24 hours prior to challenge, mice were passively immunized with a single injection of 600 mcg of murine monoclonal antibody 2H2.BE11 administered i.p. At day 0, mice were challenged with *S. aureus* MCL8538 administered i.v. The inoculum dose was 2×10^5 CFU in 100 μ l volume (Experiments 1 to 3). This low dose was found to clear spontaneously from the catheters after 4 days. Therefore, catheters were assessed for bacteria at 24 hours post challenge. At that time, mice were sacrificed and catheters harvested. The presence of bacteria on the catheters was assessed by culturing the entire catheter on TSA. If any sign of outgrowth was observed on the plate the catheter was scored as culture positive.

In two of the first three experiments, the number of culture negative catheters was significantly lower in mice passively immunized with antibody 2H2.BE11, as compared to the isotype control antibody. A fourth experiment was performed using a larger inoculum dose. In this more rigorous challenge, the dose was determined to be one in which 100% of catheters were reproducibly infected, and this infection was not spontaneously cleared by control mice (monitored over 7 days). In experiment 4, with the larger inoculum size, again, significantly fewer catheters were found to be infected in mice injected with antibody to 2H2.BE11, compared with the isotype control. Results of the four experiments are summarized in Table 8.

Table 8: Number Of Culture Negative Catheters Obtained In 4 Independent Passive Transfer Experiments Using a Murine Indwelling Catheter Model

	Number of Culture-Negative Catheters					p-value
Monoclonal	Exp#1	Exp#2	Exp#3	Exp#4	Total	
2H2.BE11	3 of 4 (75%)	6 of 8 (75%)	4 of 10 (40%)	4 of 9 (44%)	17/31 (54%)	0.0187
Isotype matched control	1 of 4 (25%)	3 of 8 (38%)	4 of 10 (40%)	0 of 9 (0%)	8/31 (25%)	

Groups of ICR mice with indwelling catheters were injected i.p. with 600 mcg of murine monoclonal antibody 24 hours prior to challenge, all monoclonals of the IgG2b isotype

5 Example 8: Ex-Vivo Pre-Opsonization of Bacteria Using anti-ORF0657n Monoclonal Antibodies
2H2.B8 (IgG1), 2H2.BE11 (IgG2b), or 13C7.IgG2b or Isotype Matched Control mAbs

To test whether monoclonal antibodies to ORF0657n are opsonic, passive protection experiments were conducted in which a lethal dose of *S. aureus* was pre-opsonized with the monoclonal antibodies 2H2.B8, 2H2.BE11, or 13C7.IgG2b, or an isotype matched control monoclonal antibody.

- 10 Pre-opsonized bacteria were then administered to mice i.p. Bacteria used in these experiments were *S. aureus* RN4220 (wild type) or RN4220.0657n. The RN4220.0657n bacteria were engineered to express ORF0657n in the absence of control by the FUR box. Therefore, they could be grown in the presence of iron and still express ORF0657n antigen on their surface. Alternatively, RN4220 (wild type) was passed 2X in a low iron medium RPMI to induce expression of 0657n on the bacteria surface.

- 15 A quantity of bacteria sufficient for 6 Balb/c mice (6 X LD₁₀₀) was incubated with 800 µg IgG at 4 °C for 1 hour, with gentle rocking. Bacteria were then pelleted and any unbound mAb removed. Antibody-opsonized bacteria were re-suspended in 2.4 mL of PBS, and 0.4 mL (1 X LD₁₀₀) was injected into each of five mice. After challenge, each inoculum was quantitated by plating on TSA to insure that equivalent CFU was given to all groups of mice and that the mAbs had not aggregated the bacteria. Survival was monitored for 3 days post challenge. Since the target antigen must be present on the surface of the bacteria for this procedure to be effective, care was taken to ensure that 0657n was expressed on the bacteria prior to opsonization. ORF0657n expression was monitored by flow cytometry using mAb 2H2.B8. The dose of opsonized bacteria injected into each mouse was 2-4 X 10⁹ CFU RN4220.0657n/mouse, or 1-2 X 10⁹ CFU RN4220(2X RPMI)/mouse.

- 25 When pre-opsonized with either 2H2.B8 or 2H2.BE11, but not an isotype matched control mAb, mice were protected from death from a lethal dose of RN4220.0657n staphylococci. The experiment was repeated twice for the IgG1 isotype and three times for the IgG2b isotype with similar results (Table 9A).

Table 9A: Ex-vivo Protection with Anti-0657n mAb

Monoclonal	Exp 1 Surviving Mice	Exp 2 Surviving Mice	Exp 3 Surviving Mice	Total
2H2.BE11 (IgG2b)	5	4	5	93% (14/15)
6G6.A8 (IgG2b)	1	0	1	13% (2/15)
PBS	1	2	0	20% (3/15)
2H2.BE11 (IgG1)	ND	4	5	90% (9/10)
10B4.H4 (IgG1)	ND	1	1	20% (2/10)

Five mice were used in each experiment. Challenge strain RN4220.0657n.pYZ119. Dose: $2-4 \times 10^9$ CFU. Test mAbs: murine anti-0657n 2H2.BE11 (IgG2b); 2H2.B8 (IgG1).

- 5 When pre-opsonized with either mAb 2H2.B8 but not an isotype matched control mAb, mice were protected from death from a lethal dose of RN4220 (2X RPMI) staphylococci. The experiment was repeated six times with similar results (Table 9B).

Table 9B: Ex-vivo Protection with Anti-0657n mAb

10

Monoclonal	# Tests	Aggregate	% Survival
2H2.B8	6	30/30	100%
10B4.IgG1 Isotype control	6	2/30	7%
13C7.IgG2b	2	0/10	0%
6G6.IgG2b Isotype control	2	0/10	0%

- 15 Murine anti-0657n 2H2 was very effective in preventing death in this lethal model. The 13C7 mAb was not effective in this model (as opposed to the previously described model illustrated in Figures 3-6). All (2H2.BE11, 2H2.B8 and 13C7.IgG2b) of the anti-0657n mAb's bind RN4220 (as demonstrated using flow cytometry) and all have opsonizing activity in the *in vitro* OPA assay. This model reflects an additional requirement for epitope specificity for enhancing survival in the peritoneum of the mouse.

Example 8: Epitope mapping studies performed with 2H2 mAb

20

The experiments described in this example provide evidence that the monoclonal antibody 2H2.BE11 recognizes a conformational epitope within ORFO657n. The experiments localized the minimal sequence within ORFO657n required for displaying the conformational epitope in a three dimensional structure recognized by 2H2 mAb. In addition, the experiments identified distinct lysine

residues within the minimal sequence of ORFO657n that become protected from reacting with small molecules when 2H2 mAb is bound to ORFO657n.

The potential ability of 2H2 mAb to recognize linear epitopes of typically 9 to 14 amino acids in length within the sequence of ORFO657n was investigated using epitope extraction and starting with an ORFO657n fragment from amino acid 42 to amino acid 486 of SEQ ID NO: 1 ("ORFO657t"). In detail: 30 ug of 2H2 mAb were immobilized by chemical cross linking to 10 mg of cyanogen bromide activated sepharose (Amersham cat. No. 17 0430 01) for each of the epitope extraction experiments. Proteolytic digests of the ORFO657t were generated with GluC (Roche Applied Science cat. No. 11 420 3997 001), Asp-N (Roche Applied Science cat. No. 11 054 589 001) or Chymotrypsin (Roche Applied Science cat. No. 11 418 467 001) and characterized by 1D/LC-MS/MS on a linear ion trap (LTQ – Thermo Electron Inc). In three individual experiments 8.4 µg of the characterized proteolytic digest from any protease was allowed to react with the immobilized antibody. Unbound peptides were washed off the antibody cross-linked beads. Potentially bound peptides were eluted with low pH and characterized by 1D/LC-MS/MS. None of the generated proteolytic peptides were recognized with high efficiency and specificity by 2H2 mAb, providing a strong indication that 2H2 mAb did not recognize a linear epitope.

The finding that 2H2 mAb did not recognize a linear sequence of ORFO657n was corroborated by a limited chemical cleavage experiment. ORFO657t was chemically cleft with CNBr for 2 hours. The resulting cleavage products were analyzed by SDS-PAGE. SDS-PAGE analysis showed 5 major bands with molecular weights of approximately 42 kDa, 35 kDa, 25 kDa, 15 kDa and 10 kDa. A Western Blot analysis with 2H2 mAb clearly showed that only the 42 kDa band was recognized by 2H2. All bands were excised from the SDS-PAGE, in-gel digest was performed, and the resulting peptides that were identified by tandem mass spectrometry were matched to corresponding sequences in ORFO657t. The result of the analysis of the major bands is shown in Table 10:

Table 10

CNBr cleavage	Binds to 2H2 mAb	ORFO657t	Calculated MW kDa
Band 42 kDa	yes	[001-356]	40.7
Band 35 kDa	no	[001-323]	36.7
Band 25 kDa	no	[001-214]	23.9
		[116-302]	21.9
Band 15 kDa	no	[215-356]	16.8
		[303-446]	16.6
Band 10 kDa	no	[114-214]	11.7
		[215-302]	10.39
		[357-446]	10.28

The importance of a fragment with a molecular weight of ~ 42 kDa was confirmed by epitope excision. In detail, 210 µg of 2H2 mAb was immobilized by chemical cross linking to 50 mg of cyanogen bromide activated sepharose (Amersham cat. No. 17 0430 01) for each of the epitope excision experiments. Then, 50 µg of intact ORF0657t was allowed to bind to the immobilized antibody and non-bound ORF0657t washed off by intensive washing with phosphate buffered saline. In three independent experiments proteases Glu-C, Trypsin and a sequential combination of GluC, AspN, Trypsin, Chymotrypsin, and Carboxy-peptidase Y were added for 5 hours or one hour per protease in the sequential combination. Peptides that were excised by the proteases during the incubation were thoroughly washed away and ORF0657t fragments that specifically bound to 2H2 mAb released with SDS loading buffer.

Fragments that specifically bound to 2H2 mAb were analyzed by SDS-page. All three of the epitope excision experiments showed exclusively one band with a molecular weight between 40 and 42 kDa in the SDS-Page analysis. Bands binding to 2H2 mAb were confirmed by Western Blot analysis. The epitope excision experiment was repeated for the Glu-C protease. This time the fragment of ORF0657t that specifically bound to 2H2 mAb was released with acidic conditions and analyzed by 1D/LC-MS/MS on a linear ion trap (LTQ, Thermo Electron). The eluted sample showed a signal (total ion count) with the expected intensity at 82-87 minutes (40%– 45% acetonitrile) and multiple charge states ($[M+67 H]^{67+}$ to $[M+30 H]^{30+}$) that deconvoluted to 42.628 kDa. A possible fragment of ORFO657t corresponding to this particular mass is sequence [012-382] of ORFO657t with a molecular weight of 42.6 kDa.

To determine which lysine residues of ORFO657t are protected from chemical reactions upon binding of 2H2 mAb, chemical labeling experiments were preformed with sulfo-NHS-acetate (Pierce Cat. No. 26777) using three different experimental conditions in the presence or absence of 2H2 mAb. See Table 11.

Table 11

Experiment	1	2	3
molar excess 2H2 mAb	0 or 3	0 or 3	0 or 3
molar excess sulfo-NHS acetate	25	500	75
Reaction temperature °C	room temperature	15	37
Reaction time	1 hour	30 minutes	2 hours

For each experiment, reaction products produced with 0 or 3 molar excess 2H2 mAb were incubated with one of three proteases resulting in 2 x 9 reaction mixtures. Experiment 1 employed

GluC, AspN and Trypsin. Experiments 2 and 3 employed GluC, AspN, and Chymotrypsin. The proteolytic peptides were then analyzed by 1D/LC-MS/MS. For each of the reactions a ratio of acetylated and non-acetylated lysine residues was calculated based on the area under curve of the total ion count (TIC) of the individual peptides. Obtained ratios were then compared between the pairs (with and without 2H2 mAb) for identical reaction conditions. A global analysis was performed for all three reaction conditions to identify lysine residues within ORF0657t that are maximally shielded upon binding of ORF0657t to 2H2 mAb. The chemical labeling experiment described above identified K76, K257 and potentially K443 as being most protected upon binding of 2H2 mAb. Protection against chemical labeling is likely due to direct binding. However, it is possible that such protection could be due to binding in close proximity to the protected sites or by long range structural changes within ORF0657nI

In summary, the above described experiments provide clear evidence that the epitope within ORF0657t that is recognized by the 2H2 mAb is conformational. The fragment of ORF0657t that is recognized by 2H2 mAb has an N-terminus located between amino acids 1 and 115 of ORF0675t and a carboxyl terminus located between amino acids 323-357 of ORF0657t. Even though it can not be excluded that protection from chemical labeling upon binding of 2H2 mAb is influenced by long range structural changes, it is very likely that areas in close proximity to Lysine 76 and Lysine 275 participate in direct antibody interaction.

Example 9: 2H2 mAb Sequence Identification

Identification of the variable light (V_L) and variable heavy (V_H) sequences of hybridoma expressed 2H2 IgG was accomplished by combining the degenerative primer PCR /overlap extension cloning process for single chain variable fragments (scFv) assembly (Krebber *et al.* *JIM 201(1)*:35-55, 1997), with high throughput screening of soluble scFv fused to a human kappa light chain constant domain or scAb material via Biacore. This allowed for fine discrimination of mutations in V_L frameworks 1, 4 and V_H frameworks 1, 4 generated by the degenerative primer method.

Briefly, RNA material was purified from the hybridoma cell line using standard methods from a Total RNA KitTM (Ambion Inc.). This material was then reverse transcribed to cDNA and utilized as template in PCR to amplify the variable regions. The conditions for the PCR amplification of the V_L and V_H chains was based upon the protocol described by Krebber *et al.* *JIM 201(1)*:35-55, 1997. The primers are designed such that a (Gly4Ser)₄ linker (SEQ ID NO: 32) is added which provides domains for a third PCR reaction in which the V_H and V_L are overlapped to create a V_L -(Gly4Ser)₄- V_H scFv.

The first set of PCR reactions to amplify the variable chains individually, were carried out in a volume of 100 μ l containing 5 μ l of the cDNA reaction, 2 μ M each of the forward and reverse primer sets for amplification of V_L and V_H , and a high fidelity PCR master mix. The reactions were denatured for 4 minutes at 94°C followed by 30 cycles of 30 seconds at 94°C, 30 seconds at 50°C, 1

minute at 72°C, and finished at a final cycle of 5 minutes at 72°C. The full length PCR products were gel purified.

To construct the full length product a third PCR reaction was done to assemble to scFv from the amplified V_H and V_L material. In a volume of 100 µl approximately 20 ng each of V_H and V_L DNA and a high fidelity PCR master mix was denatured for 5 minutes at 94°C, followed by 3 cycles of 30 seconds at 94°C, 30 s at 60°C, and 30 seconds at 72°C in the absence of primers. The modified PCR primers, SEQ ID NO: 33 and SEQ ID NO: 34 were added at a final concentration of 1 µM, and 30 cycles of 30 seconds at 94°C, 1 minutes at 60°C, and 1 minute at 72°C were performed, followed by 7 minutes at 72°C. The expected full length scFv PCR products were gel purified.

The amplified scFv material was cloned into the MP16 soluble expression vector for scAb production (Hayhurst *et al.*, *JIM* 276(1-2):185-196, 2003) and sequence analysis. A single restriction enzyme digest with SfiI was used for directional cloning into the MP16 vector. Clones with apparent full length variable heavy and variable light chains present were then expressed as scAbs in XL1-Blue cells and recovered from the periplasm using a standard osmotic shock procedure. Briefly, clones were grown at 37°C overnight in growth media containing 2% glucose and 100 µg/ml ampicillin in a 96 well format. 20 µl of the overnight culture was transferred to new media containing 0.1% glucose and 100 µg/ml ampicillin and grown until an OD₆₀₀ of 0.6 was reached. The cells were induced for scAb expression by adding IPTG at a final concentration of 0.5 mM and incubated overnight while shaking at 150 rpm, at room temperature. The scAbs were purified from the cells using a Qiagen Ni-NTA superflow robotic procedure.

To analyze each scAb periplasmic preparation for binding activity to ORF0657t, a Biacore3000 surface plasmon resonance (SPR) instrument (Uppsala, Sweden) was utilized. Standard EDC/NHS coupling was used to covalently mobilize approximately 250 resonance units of the 0657t antigen directly to the experimental flow cell surface of a CM5 sensor chip. A reference flow cell surface was activated and deactivated without coupling of protein. Each preparation was then run over the surface and association and dissociation of the scAb to antigen was measured. The surfaces were regenerated between runs by a single injection of 10 mM HCl for 20 seconds at a flow rate of 20 µl/min, followed by a 2 minute stabilization period. All samples were run in duplicate and buffer only runs were used as controls. After screening 95 clones, a clone was selected based on its binding activity. The final 2H2 clone chosen was based upon its similar affinity for ORF0657t as the original hybridoma prepared IgG material as well as comparative sequence analysis.

The amino sequence of the 2H2 V_H (SEQ ID NO: 20) and V_L (SEQ ID NO: 21) were as follows:

2H2 V_H Amino Acid Sequence (SEQ ID NO: 20)

1 DVHLVESGPG LVAPSQNLIS TCTVSGFSL RYGVHWVRQP PGKGLEWLGL
 51 IWAGGVITIYN STLMSRLSIS KDSSKSQVFL KMNSLQIDDT AIYYCAREAS
 101 RDHYFDYWGQ GTTLTVSS

2H2 V_L Amino Acid Sequence (SEQ ID NO: 21)

1 DIVMTQSPAI MSASPGEKIT MTCSASSSVS YIYWYQQKSG TSPKRWIYDT
 51 SKLASGVPFR FSGGGSGTSF SLTISSMEAE DAATYYCQOW SSNPLTFGAG
 101 TKLEIK

The underlined portions are the CDR's. CDR's were identified based on the Kabat definition. The encoding nucleic acid sequence is provided by SEQ ID NO: 24 (V_H) and SEQ ID NO: 25 (V_L).

Example 10: 2H2 IgG Chimera Expression

The variable regions for 2H2 mAb were cloned from mouse hybridoma as described in Example 9. The sequences for the variable regions were PCR amplified and DNA encoding the heavy chain variable regions were fused in-frame with DNA encoding the IgG1 constant region whereas DNA encoding the light chain variable region were fused in-frame with DNA encoding the kappa constant region. The cloning procedure for the resulting antibody expression vectors is described below.

The variable regions were PCR amplified. PCR reactions were carried out in a volume of 25 µl containing high fidelity PCR master mix, template volume 1 µl and forward and reverse primers: 1 µl each. PCR condition was 1 cycle of 94°C, 2 minutes, 25 cycles of 94°C, 1.5 minutes; 60°C, 1.5 minutes; 72°C, 1.5 minutes and 72°C, 7 minutes; 4°C until removed and cloned in-frame with leader sequence at the 5'-end and constant region at the 3'-end using In-Fusion strategy. The following primers were used: Light chain forward, 5'- ACAGATGCCAGATGCGATATTGTGATGACCCAGTCT (SEQ ID NO: 28); Light chain reverse, 5'- TGCAGCCACCGTACGTTTTATTTCCAGCTTGGTCCC (SEQ ID NO: 29); Heavy chain forward, 5'- ACAGGTGTCCACTCGGATGTGCACCTGGTGGAGTCA (SEQ ID NO: 30); and Heavy chain reverse, 5'- GCCCTTGGTGGATGCCGAGGAGACTGTGAGAGTGGT (SEQ ID NO: 31). The DNA sequences for all the clones were confirmed by sequencing.

The amino acid sequences deduced from DNA sequences are:

Mouse 2H2 Variable and Human Kappa Constant Region Amino Acid Sequence (SEQ ID NO: 22)

1 DIVMTQSPAI MSASPGEKIT MTCSASSSVS YIYWYQQKSG TSPKRWIYDT
 51 SKLASGVPFR FSGGGSGTSF SLTISSMEAE DAATYYCQOW SSNPLTFGAG
 101 TKLEIKRTVA APSVFIFPPS DEQLKSGTAS VVCLLNNFYP REAKVQWKVD
 151 NALQSGNSQE SVTEQDSKDS TYSLSSTLTL SKADYEKHKV YACEVTHQGL
 201 SSPVTKSFNR GEC

Mouse 2H2 Variable and Human IgG1 Constant Region Amino Acid Sequence (SEQ ID NO: 23)

1 DVHLVESGPG LVAPSQNLSI TCTVSGFSLS RYGVHWVRQP PGKGLEWLGL
 51 IWAGGVTIYN STLMSRLSIS KDSSKSQVFL KMNSLQIDDT AIYYCAREAS
 5 101 RDHYFDYWQ GTTLTVSSAS TKGPSVFPLA PSSKSTSGGT AALGCLVKDY
 151 FPEPVTVSWN SGALTSGVHT FPAVLQSSGL YSLSSVVTVP SSSLGTQTYI
 201 CNVNHKPSNT KVDKRVEPKS CDKTHTCPPC PAPELLGGPS VFLFPPKPKD
 251 TLMISRTPEV TCVVVDVSHE DPEVKFNWYV DGVEVHNAKT KPREEQYNST
 301 YRVVSVLTVL HQDWLNGKEY KCKVSNKALP APIEKTISKA KGQPREPQVY
 10 351 TLPPSREEMT KNQVSLTCLV KGFYPSDIAV EWESNGQPEN NYKTTTPVLD
 401 SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH EALHNHYTQK SLSLSPGK

The variable regions are underlined.

The antibodies were expressed in 293EBNA monolayer cells. The plasmids were transfected using PEI based transfection reagents. The transfected cells were incubated in Opti-MEM serum free medium and the secreted antibodies were purified from medium using protein A/G affinity chromatography. The concentration of purified antibodies was determined by OD280 nm and the purity was measured by LabChip™ capillary electrophoresis.

The expression of both light and heavy chains was driven by human CMV promoter and bovine growth hormone polyadenylation signal. (Shiver *et al.*, *Ann. N.Y. Acad. Sci.*, 772:198-208, 1995.) The leader sequence in the front mediated the secretion of antibodies into the culture medium. The leader sequence for the heavy chain was MEWSWVFLFFLSVTTGVHS (SEQ ID NO: 26) and for the light chain was MSVPTQVLGLLLWLTDAARC (SEQ ID NO: 27). The expression vectors carry oriP from EBV viral genome for prolonged expression in 293EBNA cells and the bacterial sequences for kanamycin selection marker and replication origin in *E. coli*.

The antibodies were expressed in 293EBNA monolayer cells. The plasmids were transfected using PEI based transfection reagents. The transfected cells were incubated in Opti-MEM serum free medium and the secreted antibodies were purified from medium using protein A/G affinity chromatography. The concentration of purified antibodies was determined by OD280nm and the purity by LabChip capillary electrophoresis.

30

Example 11: Affinity Determination

Comparative analysis was performed on 2H2 mAb as hybridoma material, scAb and a chimeric antibody. 2H2 mAb V_H and V_L region were cloned and expressed as an IgG chimera as described in Example 10. scAb was cloned into the MP16 vector (Example 9), which produces a scFv with a Human Kappa chain tag fused to it. As further described below, the antigen affinity was not significantly different among the constructs.

To measure a 1:1 interaction between the binding domain and the antigen, the experimental set up on Biacore was modified depending on whether antibody fragment or full length IgG

was analyzed. For IgG measurements, the IgG was captured to the surface as ligand and ORF0657t was run as analyte. For antibody fragment analysis, ORF0657t was bound to the surface and the antibody fragment was run as the analyte. This demonstrated that the affinity of the original 2H2 mAb hybridoma material to the ORF0657t antigen shows no significant change upon recombinant cloning (Table 12).

- 5 Data were acquired via surface plasmon resonance on a Biacore 3000; each analyte was run at multiple concentrations, with two replicates per concentration. Data were analyzed with BIAevaluation (Biacore, Inc.) with simultaneous fits of entire concentration series. Fit parameters are listed in Table 12.

Table 12

	On-rate k_a (1/Ms)	Off-rate k_d (1/s)	Affinity, KD	χ^2 global fit
2H2 murine IgG2b	6.10 E+04	2.01 E-03	33nM	0.902
2H2 scAb	4.91 E+04	1.91 E-03	39nM	0.429
10 2H2 IgG chimera	1.10 E+05	2.73 E-03	25nM	0.295

Example 12: ORF0657n Based Sequences

The highlighted amino acids (indicated by bold and underlying) present in SEQ ID NOs:

- 15 4-19 show amino acid alterations to ORF0657n:

0657n (SEQ ID NO: 1)

20 MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKAVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
 MKKENGEQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTPVKDYAYIRFSVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQRVRTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKAFTKANTDKS
 25 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKT PATKPTKGEVESSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

0657nC/e (SEQ ID NO: 2)

30 MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKAVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
 MKKENGEQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTPVKDYAYIRFSVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 35 TTNDYWKDFMVEGQRVRTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKT PATKPTKGEVESSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKNLEHHHHH

0657nH/y (SEQ ID NO: 3)

MAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVANAVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKA
 TNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFEMKKKGDTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQF
 WRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGTKAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEE
 5 DYKAEKLLAPYKKA~~KT~~LERQVYELN~~KI~~QDKLPEKLKAEYKKKLED~~TK~~KALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDT
 KYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMETTNDDYWKDFMVEGQVRV~~TI~~SKDAKN~~NR~~TIIFPYVEGKT
 LYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKSNKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDD
 NKQLPSVEKENDASSES~~SG~~KDKTPATKPTKGEVSSSTPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSA
 PLQKANIKNTNDGHTQSQNNKNTQENKAKS

10

SEQ ID NO: 4

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
 MKKKGDTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
 15 ~~KE~~VKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKA~~KT~~LERQVYELN~~KI~~QDKLPEKLKA
 EYKKKLED~~TK~~KALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMME
 TTNDDYWKDFMVEGQVRV~~TI~~SKDAKN~~NR~~TIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSES~~SG~~KDKTPATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 20 SNKDMTLPLMALLALSSIVAFVLPKRKKN

SEQ ID NO: 5

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPN~~WR~~PIDFE
 25 MKKKGDTQQFYHYASSV~~E~~PARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
~~KE~~VKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKA~~KT~~LERQVYELN~~KI~~QDKLPEKLKA
 EYKKKLED~~TK~~KALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMME
 TTNDDYWKDFMVEGQVRV~~TI~~SKDAKN~~NR~~TIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSES~~SG~~KDKTPATKPTKGEVSSST
 30 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPLMALLALSSIVAFVLPKRKKN

SEQ ID NO: 6

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 35 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPN~~WR~~PIDFE
 MKKKGDTQQFYHYASSV~~E~~PARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKA~~KT~~LERQVYEL~~E~~KIQDKLPEKLKA
 EYKKKLED~~TK~~KALDEQVKS~~AI~~TEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMME
 TTNDDYWKDFMVEGQVRV~~TI~~SKDAKN~~NR~~TIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 40 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSES~~SG~~KDKTPATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPLMALLALSSIVAFVLPKRKKN

SEQ ID NO: 7

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 5 MKKKDGTQQFYHYASSVEPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLEQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALAEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVESSST
 10 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 8

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 15 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLEQVYELEKIQDKLPEKLKA
 EYKKKLEDTKKALAEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 20 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVESSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 9

25 MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTKSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTQKDYAYIRFSVSNGT
 KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLEQVYELEKIQDKLPEKLKA
 EYKKKLEDTKKALAEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 30 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVESSST
 TPTKVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPMLALLALSSIVAFVLPKRKN

35 SEQ ID NO: 10

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
 MKKKDGTQQFYHYASSVEPARVIFTKSKPEIELGLQSGSTWRKFEVYEGDKKLPIKLVSYDTQKDYAYIRFSVSNGT
 40 KEVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLEQVYELEKIQDKLPEKLKA
 EYKKKLEDTKKALAEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVESSST

TPTKVVSTTQNVAKPTTASSKTTKD VVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 11

5 MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
MKNDKGTQQFYHYASSVEPARVIFTKSKPIIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTDKDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLERQVYELEKI QDKLPEKLKA
EYKKKLEDTKKALAEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE
10 TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKT PATKPTKGEVSSST
TPTKVVSTTQNVAKPTTASSKTTKD VVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPKRKN

15 SEQ ID NO: 12

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
MKNDKGTQQFYHYASSVEPARVIFTKSKPIIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTDKDYAYIRFSVSNGT
KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLERQVYELEKI QDKLPEKLKA
20 EYKKKLEQTKKALAEQVKS AITEFQNVQPTNEKMTDLQDAHYVVYESVENSESMDTFVKHPIKTGMLNGKKYVMVE
TTNDYWKDFMVEGQVRVTISKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKT PATKPTKGEVSSST
TPTKVVSTTQNVAKPTTASSKTTKD VVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
SNKDMTLPMLALLALSSIVAFVLPKRKN

25

SEQ ID NO: 13

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNWRPIDFE
MKNDKGTQQFYHYASSVEPARVIFTKSKPIIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTDKDYAYIRFSVSNGT
30 KEVKIVSSTHFNNKEEKYDYTLMVFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLERQVYELEKI QDKLPEKLKA
EYKKKLEQTKKALAEQVKS AITEFQNVQPTNEKMTDLQDAHYVVYESVENSESMDTFVKHPIKTGMLNGKKYVMVE
TTNDYWKDFMVEGKVRVTISKDAKNNTRTIIFPYVEGKALYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKT PATKPTKGEVSSST
TPTKVVSTTQNVAKPTTASSKTTKD VVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
35 SNKDMTLPMLALLALSSIVAFVLPKRKN

SEQ ID NO: 14

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTTEAQPKEAVASPTTTSEKAPETKPVAN
AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIKDKDHSAPNSRPIDFE
40 MKKKDGTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVKDYAYIRFSVSNGT
KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKAKTLERQVYELNKLQEKLPEKLKA
EYKKKLEDTKKALDEQVKS AITEFQNVQPTNDKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVMVE

TTNDDYWKDFMVEGQSVRTISKDAKNNTRTIIFFPYIEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKTPATKPTKGEVESSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPLMALLALSSIVAFVLPRKRKN

5

SEQ ID NO: 15

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNNTYPILNQELRDAIKNPAIKDKHSA PNSRPIDFE
 MKKKDGTQQFYHYASTVKPARVIFTDTKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDSVKDYAYIRFSVSNGT
 10 RAVKIVSSTHYNNKEEKYDYTLMEFAQPIYNSADKYKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
 EYKKKLDTKKALDDQVKSAYTEFQNVQPTNEKMTDLQDTKYVVFESVENNESVMDTFVKHPIKTGMLNGKKYVME
 TTNDDYWKDFIVEGQVRVTISKDAKNNTRTIIFFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKTPATKPTKGEVESSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 15 SNKDMTLPLMALLALSSIVAFVLPRKRKN

SEQ ID NO: 16

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNNTYPILNQELREA IKNPAIIDKHSA PNSRPIDFE
 20 MKKKDGTQQFYHYASSVKPARVIFTDSGPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYVME
 TTNDDYWKDFMVEGQVRVTISKDAKNNTRTIIFFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKTPATKPTKGEVESSST
 25 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPLMALLALSSIVAFVLPRKRKN

SEQ ID NO: 17

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 30 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNNTYPILNQELREA IKNPAIIDKHSA PNSRPIDFE
 MKKKDGTQQFYHYASSVKPARVIFTDSGPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFVSNGT
 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKS AITEFQNVQPTNEKMTDLQDTKYVVYESEENNESMMDTFVKHPIXTGMLNGKKYVME
 TTNDDYWKDFMVEGQVRVTISKDAKNNTRTIIFFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 35 NKKEQQDNSAKKEATPATPSKPTSPVEKESQKQDSQKDDNKQLPSVEKENDASSESGKDKTPATKPTKGEVESSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 SNKDMTLPLMALLALSSIVAFVLPRKRKN

SEQ ID NO: 18

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREAIKNPAIDDKHSAPNSRPIDFE
 MKKKDGTQQFYHYASSVKPARVIFTDSQPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFPVSNGT
 5 KAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKDEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEKLKA
 EYKKKLEDTKKALDEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESEENNESMMDTFVKHPIYTGMLNGKKYVMVE
 TTNDYWKDFMVEGQVRVTSKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDKEAFTKANTDKS
 NKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEVSSST
 TPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLPQTGEE
 10 SNKDMTLPLMALLALSSIVAFVLPRKRKN

SEQ ID NO: 19

MNKQQKEFKSFYSIRKSSLGVASVAISTLLLLMSNGEAQAAAEETGGTNTEAQPKEAVASPTTTSEKAPETKPVAN
 AVSVSNKEVEAPTSETKEAKEVKEVKAPKETKEVKPAAKATNNTYPILNQELREGSEAIKNPAIKDKHSAPNSRPI
 15 DFEMKKDGTQQFYHYASSVKPARVIFTDSKPEIELGLQSGQFWRKFEVYEGDKKLPIKLVSYDTVVDYAYIRFSVS
 NGTKAVKIVSSTHFNNKEEKYDYTLMEFAQPIYNSADKFKTEEDYKAEKLLAPYKKAKTLERQVYELNKIQDKLPEK
 LKAEYKKKLEDTKKALDEQVKSATEFQNVQPTNEKMTDLQDTKYVVYESVENNESMMDTFVKHPIKTGMLNGKKYM
 VMETTNDYWKDFMVEGQVRVTSKDAKNNTRTIIFPYVEGKTLYDAIVKVHVKTIDYDGQYHVRIVDVDKEAFTKA
 NTDKSNKKEQQDNSAKKEATPATPSKPTPSPVEKESQKQDSQKDDNKQLPSVEKENDASSESQKDKTPATKPTKGEV
 20 ESSSTTPTKVVSTTQNVAKPTTASSKTTKDVVQTSAGSSEAKDSAPLQKANIKNNTNDGHTQSQNNKNTQENKAKSLP
 QTGEESNKDMTLPLMALLALSSIVAFVLPRKRKN

Other embodiments are within the following claims. While several embodiments have
 been shown and described, various modifications may be made without departing from the spirit and
 25 scope of the present invention.

WHAT IS CLAIMED IS:

1. An isolated antigen binding protein comprising a first variable region and a second variable region, wherein said binding protein binds to a target region selected from the group consisting of: mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target region.
2. The binding protein of claim 1, wherein said target region is the mAb 2H2.BE11 target region and said first variable region is a V_h region comprising:
 - a first V_h CDR comprising amino acids 36-45 of SEQ ID NO: 20 or a sequence differing from amino acids 36-45 by one amino acid;
 - a second V_h CDR comprising amino acids 50-65 of SEQ ID NO: 20 or a sequence differing from amino acids 50-65 by one amino acid; and
 - a third V_h CDR comprising amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.
3. The binding protein of claim 2, wherein said second variable region is a V_l region comprising:
 - a first V_l CDR comprising amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid;
 - a second V_l CDR comprising amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and
 - a third V_l CDR comprising amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.
4. The binding protein of claim 3, wherein said binding protein is an antibody.
5. The binding protein of claim 4, wherein said antibody is a monoclonal antibody.
6. The binding protein of claim 4, wherein said V_h region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and said V_l region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.
7. The binding protein of claim 6, wherein said binding protein is an antibody comprising (a) a heavy chain comprising said V_h region, and a human hinge, CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄ subtype; and (b) a light chain comprising said V_l region, and either a human kappa C_L or human lambda C_L.

8. The binding protein of claim 3, wherein
said V_h region comprises said first V_h CDR consisting of amino acids 36-45 of SEQ ID
NO: 20, said second V_h CDR consisting of amino acids 50-65 of SEQ ID NO: 20, and said third V_h
5 CDR consisting of amino acids 98-107 of SEQ ID NO: 20 and;
said first V_l region comprises said first V_l CDR consisting of amino acids 24-33 of SEQ
ID NO: 21, said second V_l CDR consisting of amino acids 49-55 of SEQ ID NO: 21, and said third V_l
CDR consisting of amino acids 88-96 of SEQ ID NO: 21.
- 10 9. The binding protein of claim 8, wherein said binding protein is an antibody.
10. The binding protein of claim 9, wherein said antibody is a monoclonal antibody.
11. The binding protein of claim 9, wherein said V_h region is either SEQ ID NO: 20,
15 a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20; and said V_l region is either SEQ ID
NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.
12. The binding protein of claim 8, wherein said binding protein is an antibody
comprising (a) a heavy chain comprising said V_h region, and a human hinge, CH_1 , CH_2 , and CH_3
20 regions from an IgG1, IgG2, IgG3 or IgG4 subtype; and (b) a light chain comprising said V_l region, and
either a human kappa C_L or human lambda C_L .
13. The binding protein of claim 12, wherein said heavy chain consists essentially of
the amino acid sequence of SEQ ID NO: 22; and said light chain consists essentially of the amino acid
25 sequence of SEQ ID NO: 23.
14. A nucleic acid comprising a recombinant gene comprising a nucleotide sequence
encoding an antibody variable region that binds to a target region selected from the group consisting of:
mAb IG3.BD4 target region, mAb 2H2.BE11 target region, mAb 13C7.BC1, and mAb 13G11.BF3 target
30 region.
15. The nucleic acid of claim 14, wherein said target region is the mAb 2H2.BE11
target region and said variable region is a V_h region comprising:
a first V_h CDR comprising amino acids 36-45 of SEQ ID NO: 20 or a sequence differing
35 from amino acids 36-45 by one amino acid;
a second V_h CDR comprising amino acids 50-65 of SEQ ID NO: 20 or a sequence
differing from amino acids 50-65 by one amino acids; and

a third V_H CDR comprising amino acids 98-107 of SEQ ID NO: 20 or a sequence differing from amino acids 98-107 by one amino acid.

16. The nucleic acid of claim 15, wherein said V_H region comprises said first V_H CDR consisting of amino acids 36-45 of SEQ ID NO: 20; said second V_H CDR consisting of amino acids 50-65 of SEQ ID NO: 20; and said third V_H CDR consisting of amino acids 98-107 of SEQ ID NO: 20.

17. The nucleic acid of claim 14, wherein said variable region is either SEQ ID NO: 20, a humanized SEQ ID NO: 20, or a de-immunized SEQ ID NO: 20.

18. The nucleic acid of claim 17, wherein said recombinant gene encodes an antibody heavy chain comprising said variable region, a human hinge, and CH₁, CH₂, and CH₃ regions from an IgG₁, IgG₂, IgG₃ or IgG₄ subtype.

19. The nucleic acid of claim 14, wherein said variable region is a V_L region comprising:
a first V_L CDR comprising amino acids 24-33 of SEQ ID NO: 21 or a sequence differing from amino acids 24-33 by one amino acid;
a second V_L CDR comprising amino acids 49-55 of SEQ ID NO: 21 or a sequence differing from amino acids 49-55 by one amino acid; and
a third V_L CDR comprising amino acids 88-96 of SEQ ID NO: 21 or a sequence differing from amino acids 88-96 by one amino acid.

20. The nucleic acid of claim 19, wherein said first V_L region comprises said first V_L CDR consisting of amino acids 24-33 of SEQ ID NO: 21, said second V_L CDR consisting of amino acids 49-55 of SEQ ID NO: 21, and said third V_L CDR consisting of amino acids 88-96 of SEQ ID NO: 21.

21. The nucleic acid of claim 14, wherein said variable region is either SEQ ID NO: 21, a humanized SEQ ID NO: 21, or a de-immunized SEQ ID NO: 21.

22. The nucleic acid of claim 17, wherein said recombinant gene encodes an antibody light chain comprising said variable region and a human kappa or lambda C_L.

23. A recombinant cell comprising one or more nucleic acids of any one of claims 14-22.

24. The recombinant cell of claim 23, wherein said cell comprises both the nucleic acid of claim 18 and the nucleic acid of claim 22.

25. A method of producing protein comprising an antibody variable region
5 comprising the steps of:
a) growing the recombinant cell of claim 23 under conditions wherein said protein is expressed; and
b) purifying said protein.

10 26. A method of producing protein comprising an antibody variable region comprising the steps of:
a) growing the recombinant cell of claim 24 under conditions wherein said protein is expressed; and
b) purifying said protein.

15 27. A pharmaceutical composition comprising the binding protein of any one of claims 1-13 and a pharmaceutically acceptable carrier.

20 28. A method of detecting the presence of an OFR0657n antigen in a solution or on a cell comprising the steps of: (a) providing the binding protein of any one of claims 1-13 to said solution or said cell; and (b) measuring the ability of said binding protein to bind to said antigen present in said solution or to said cell.

25 29. A method of treating against an *S. aureus* infection in a patient comprising the step of administering to said patient an effective amount of the binding protein of any one of claims 1-13.

30 30. The method of claim 29, wherein said antigen binding protein is administered in conjunction with surgery or a foreign body implant.

31. A cell line producing a protein that is either mAb 1G3.B3, mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11, or that competes with either mAb 1G3.B3., mAb 2H2.B8, mAb 13C7.D12, or mAb 13G11.C11 for binding to ORF0657n.

32. The cell line of claim 31, wherein said line is either ATCC No: PTA-7124,
35 ATCC No: PTA-7125, ATCC No: PTA-7126 or ATCC No: PTA-7127.

1/5





FIG. 1

2/5

**FIG. 2**

3/5




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4A

4/5




FIG. 4B




FIG. 5A





FIG. 5B

5/5

**FIG. 5C**